



The Biology Curator




The Publication of the Biology Curator's Group

ISSUE 10 Special Supplement, Funded by the Natural History Museum NOVEMBER 1997

A REVIEW OF TECHNIQUES USED IN THE PREPARATION, CURATION AND CONSERVATION OF MICROSCOPE SLIDES AT THE NATURAL HISTORY MUSEUM, LONDON

PAUL A. BROWN
The Natural History Museum



Abstract

Permanent microscope slide mounts have been an integral part of the Natural History Museum's (NHM) collections since microscopes were first used in the study of natural history specimens. This paper discusses the methods, materials and mountants used in the preparation of such slides and their storage based on the findings of a museum-wide survey. Over the years, many recipes for slide mounting media have been tried, the mountants differing in their optical properties and their ageing characteristics. Using the collections as a data base of deterioration dating back to the early 19th century, conclusions are drawn as to which are the most suitable mounting media and which are the most suitable conservation techniques and housing methods. Appended are a list of recipes and a full bibliography.

Contents

	<i>Page No.</i>		<i>Page No.</i>
		Abstract	
1.	3	Introduction	
2.	3	Microscope slide making and curatorial practices.	
2.1		Microscope slides	
2.2		Dry mounts	
2.3		The mounting medium	
2.4		Coverslip supports	
2.5		Liquid mounts	
2.6		Alternatives to microscope slide mounts	
2.7		Chromosome preparation slides	
2.8		Electron microscopy	
2.9		Methods for specimen preparation prior to slide mounting	
2.9.1		Maceration	
2.9.2		Tissue embedding and sectioning	
2.9.3		Stains and bleaching agents (see end of Appendix 3)	
2.10		Microscope slide mounting methods	
2.11		Ringing the coverslip	
2.12		Labels	
2.13		Immersion oils	
2.14		Storage	
3.	6	The choice of microscope slide mountants, slide mounting media used at the NHM	
3.1		Refractive indices (see Appendix 2.)	
3.2		Natural resinous media	
3.2.1		Canada balsam	
3.2.2		Phenol balsam	
3.2.3		Euparal	
3.3		Synthetic resinous media (plastics)	
3.3.1		Cellofas = Carboxy Methyl Cellulose	
3.3.2		DePeX = Distrene plasticiser xylene	
3.3.3		Dimethyl hydantoin formaldehyde (DHFM)	
3.3.4		Naphrax	
3.3.5		Permout	
3.3.6		Histomount	
3.3.7		PolyVinyl Lactophenol (PVLPL)	
3.4		Gum chloral mountants	
3.4.1		Crystallisation	
3.4.2		Blackening	
3.5		Liquid mounts	
3.5.1		Glycerine-glycerol	
3.5.2		Karo corn syrup	
3.5.3		Lactophenol	
3.6		Ringing media	
3.7		Adhesives	
4.		Literature survey	11
5.		Conclusion. Which are the best mountants and what future do slide collections have	11
6.		Acknowledgements	12
7.		Appendix 1: Mountant recipes	12
8.		Appendix 2: Refractive indices	18
9.		Appendix 3: Materials & Suppliers	19
10.		Bibliography	19
		PLATES	27
		Curriculum vitae	Inside Back Cover

1. Introduction

Microscopes were rare and little used until the first half of the nineteenth century, when rapid advances in optical research and in manufacturing methods produced cheaper and more reliable instruments. These microscopes became available to the serious biologist and geologist (Allen, 1976: 128) and their availability is reflected in the early 19th century dates for some microscope slides within the NHM's collections. The new microscope slide mounts rapidly developed into collections in the possession of private individuals, universities, hospitals and other institutions. Collections were made of whole organisms as well as sections of minerals, plant stems, worms etc and of microscopic parts of larger organisms such as the cells, pollen grains, spores, hairs, scales, insect genitalia and diseased tissue. Some of this material opened up new avenues for taxonomic research. As their taxonomic value grew, slide preparations increasingly featured in natural history museum collections. In addition to slides made by museum staff, private collections were also transferred to museums for safe keeping, particularly from institutions which no longer had the resources to properly care for such collections.

Deterioration of some mounting media was noted within the NHM aphid (Hemiptera) collection and a programme of survey and rescue of important material was put into operation by the author. As the subject of an MA dissertation, a museum wide study was undertaken to ascertain the condition of other slide collections. Such collections have been thought less prone to deterioration than spirit preserved or dry material and conservation problems relating to them have received little attention. This paper is a revised version of the MA project study.

2. Microscope slide making and curatorial practices.

2.1 Microscope slides.

Initially microscope preparations were made on many different sizes and shapes of glass 'slides'. This can still present a problem in storage geared for the standard and now universally accepted, sized slide which is 75 x 25 x 0.8-1.2 mm. In the examination of most microscope slides it is desirable for transmitted light to pass through the slide, so the slide, mountant (supporting and preserving medium) and coverslip are usually transparent. Coverslips can be square or round, although the latter are preferred as the corners on a square coverslip can catch, causing it to split. Some glass slides can have a shallow cavity ground in the slide to facilitate the mounting of thicker specimens. The higher the magnification used in the examination of the specimen, the thinner should be the coverslip (the thickness varying between 0.085 to 0.35 mm) and the thinner should be the depth of mountant. In addition to glass, slides may be of metal, plastic or wood with a central hole and some dry mounts rely on reflected light so that the slides can be totally opaque. The Cobb slide is an aluminium strip with a hole drilled centrally and is flanged along the long edges. The flange holds card squares with documentation and the specimens are mounted between two coverslips positioned over the central hole (See Fig. 1) (Westheide & Pursche, 1988: 153). The Higgins-Shirayama slide is a further

development made of plastic. Two sheets of plastic, one with a 16 mm wide hole and the other with an 18 mm wide hole, are fused together so that an 18 mm diameter coverslip can be fitted and stuck to the 2 mm wide shelf (Westheide & Purschke, 1988: 153). Similar slides made of perspex are found in the Rothschild insect collection (Fig. 2). The specimen is then arranged in the mountant and another smaller coverslip placed on top so that high magnification viewing is possible from both sides of the slide, unlike a glass slide mount where the thickness of glass on the reverse side hampers or prevents close inspection. High magnification of thick mounts is limited as it is not possible for the microscope objective to be placed close enough to the subject.

2.2 Dry mounts.

Dry mounts are used when a preservative is not required, for example microlepidoptera wings (Fig. 3), and for small opaque specimens such as copepod shells which are viewed using reflected light. These slides need not be transparent so glass can be replaced by stronger opaque materials. Humes & Gooding (1964: 238) discuss the use of wooden slides for copepods. Ostracods are often mounted on black card on wood, cardboard, glass or plastic slides as dry mounts (Fig. 4). The black and white plastic slides figured (Fig. 5) have a clip-on and removable transparent square coverslip to protect the specimens from dust. The dust problem is described by Green for micropalaeontological specimens (1995: 162).

The specimens are attached with spots of adhesive which must not obscure the specimen, and protected in a cavity slide or by solid rings beneath the coverslip. As the coverslip is not supported from below by mountant any pressure exerted from above can cause the glass coverslip to break, exposing the specimen to damage. Such damage can also occur when liquid mounts dry out (due to evaporation through cracked ringing media for example) because the liquid also supports the coverslip. Wooden and card slides with wells cut in them are suitable for housing electron microscopy stubs as found in the Zoology Department of the NHM (Fig. 6).

2.3 The Mounting Medium.

Mounting media may be liquid, gum or resinous, soluble in water, alcohol or other solvents and be sealed from the external atmosphere effect by non-soluble ringing media. The refractive index of the mountant should be chosen to be either similar to, or contrast with, the specimen, depending on the method of viewing of the slide and the nature of the specimen.

2.4 Coverslip supports.

There are a number of methods to support the cover slip when the mount is thick. The support can take the form of a glass or metal ring, cut sections of insect pins, or glass beads or celluloid (Fig. 7) placed around the specimen so as to reduce flattening of the specimen by the coverslip. Cardboard rings (as described by Jobling (1938: 55)) can be cut to the right thickness for the mounted specimens, as used for Euparal mounts of small Hymenoptera, but are not considered suitable by Gray (1954). Problems arise when the

solvent evaporates and the coverslip can be broken as stresses build, so extra mountant should be added at a later date to prevent this. Cavity slides also have this problem with the coverslip distorting in the centre. Celluloid cut into 1 mm. squares is a good support as it 'gives' slightly as the mountant shrinks. Thick square card labels or strips or beads (Fig. 8) can be fixed on each side of the coverslip to act as 'spacers' to protect the mount.

2.5 Liquid mounts.

Liquid mounts require leak proof cells to contain and seal in the liquid. These can be made of card, metal, glass or a suitable water or solvent resistant ringing medium or cement with further layers of soft paraffin wax if required. Such cells can also be used for dry mounts.

2.6 Alternatives to microscope slide mounts.

Most slide mounts are essentially two dimensional with specimens usually flattened dorso-ventrally but not so flat that the difference between the dorsal and ventral surfaces cannot be detected. Some workers dissect specimens and place different parts of the specimen under different small coverslips on the same slide (Fig. 9). As an alternative to slide mounting, associated parts of larger organisms can be placed in vials in liquid preservative to preserve their three dimensional character. The NHM's collection of Butterfly genitalia are placed in vials of glycerine and pinned through the cork on the mounted insect pins. Robinson (1976: 129) considers this technique to be "abhorrent", stressing the likelihood of damage and loss in handling. Vials of liquid preservative may dry out and thus require frequent inspection. For this reason, slide mounts are preferable. Coleoptera genitalia (Cooter, 1991) or Diptera genitalia (Coe, 1966: 19) have been mounted in slide mountant (Euparal and Canada balsam) on small squares of celluloid (cellulose acetate) or card, and also placed on the insect pin (Imms, 1929: 166). Robinson (1976: 129) also condemns this method as "obnoxious" and suggests that genitalia preparations should always be on microscope slides. A further method previously used was to mount the specimen between two coverslips over a hole in card (Walker & Crosby, 1988: 21), sealed with glued paper which can then be pinned or enveloped (Fig. 10). The latter two methods are no longer practised in the NHM Entomology department.

2.7 Chromosome preparation slides

Chromosome preparation slides can be made from and associated with specimens before preservation of the latter in the main collection. The NHM Aphidoidea chromosome collection numbers some 5,000 slide mounts in DePeX (which are unfortunately deteriorating). A similar collection of simuliid fly chromosome slides mounted in Euparal are also deteriorating but at a slower rate. This deterioration is probably due to the ephemeral nature of chromosomes rather than that of the mountants. Both these collections are considered to be secondary in importance to the collection of photographs of the chromosome preparations made when fresh. The NHM's 10,000 plant chromosome slides are also mounted in Euparal.

2.8 Electron microscopy

Electron microscopy (EM) is increasingly being used in taxonomy, demanding many new preparation techniques and mountants. However, because such preparations are not normally in the form of permanent glass microscope slides, they are not covered in this paper. Some embedding media used in electron microscopy such as polymer resins, as discussed by Smith & Tyler (1984: 260) are suitable for conventional light microscopy. Few new mounting media are now being developed for conventional microscopy as most development is now within the electron microscopy field. The curation of scanning electron microscope stubs is discussed by Julia Golden (1989: 17-26). Stephen Russell (1989) documents a technique for reconditioning SEM treated diatoms for conventional slide mounting. The copepod collection houses SEM stubs in card well slides (Fig 6).

2.9 Methods for specimen preparation prior to slide mounting.

2.9.1 Maceration.

Before specimens of insects and some other groups can be made into permanent slide mounts, they must be prepared or 'macerated' using the appropriate method for the chosen mountant. Body tissues, fat, secretions and wax often need to be cleared or denatured, making the organism translucent to facilitate the examination of the surface or exoskeletal structures of the organism. The body contents must not be fixed with formaldehyde which preserves the body tissues so that they then cannot be cleared. Sodium hydroxide or potassium hydroxide are commonly used chemical macerating agents, at varying strengths, for varying periods, and at different temperatures depending on the size and fragility of the specimens. When specimens preserved in aqueous solutions are destined for resinous mounts, they require dehydration as many of the natural and synthetic resin mountants do not mix with water. Dehydration is achieved by placing the specimens progressively through a series of aqueous ethyl alcohol solutions, 50%, 80%, 95% until 100% ethanol is reached. Alternatively, two rinses in glacial acetic acid may suffice (suitable for aphids but not Thysanoptera). Preparation must be done in such a way that the fluid in the specimen is compatible with the mountant thus avoiding opaqueness or osmotic collapse and distortion, which renders the specimen useless for taxonomic study.

2.9.2 Tissue embedding and sectioning.

Thin tissue sections are cut after the tissue is embedded in a medium which holds the tissue together. To make a permanent slide preparation of the section it has to be glued to the slide with a suitable adhesive which should have a similar refractive index to the glass of the slide (RI = 1.51), to the embedding medium and to the mounting medium. Cellulose nitrate has a refractive index of 1.49-1.51 and polyisobutylene adhesive has a refractive index of 1.50-1.51 (Fink, 1987: 97) With possibly three different media involved in the preparation of a plant section one must make sure that they are compatible, both chemically and optically, as the boundary between the media could reduce the visibility of the section. Rawlins (1992: 53) discusses the use

of embedding media and glass microtome or vibratome cutting tools.

2.9.3 Stains and bleaching agents (see also Appendix 3).

For very translucent specimens, a stain may be required to improve the visibility of the specimen before it is placed into the mountant. There are many stains available which stain different constituent chemical parts of the organism. One must choose the appropriate stain for both the organism and the slide mountant. New (1974: 23) suggests that some stains deteriorate in certain mountants. He suggests that Acid Fuchsin, Fast green and Lignin pink can be fixed in Euparal and Canada balsam, but that Chlorazol Black E is better with polyvinyl lactophenol or methyl cellulose/Carbowax media. This is not the experience of the Microlepidoptera section at the NHM which regularly uses Chlorazol Black E with Euparal and has witnessed little or no deterioration over thirty years. In the NHM Coccoidea collection Acid Fuchsin is used with Canada balsam. Some fading of this stain does occur and remnants of clove oil, the final clearing fluid may be involved as Acid Fuchsin stained specimens stored in clove oil fade within a few days (J. Martin pers. comm.). When using phase contrast, the stain is considered as useful only to locate the organism on the expanse of the slide. It is widely accepted that acid stains will fade in neutral or alkaline mountants but that neutral mountants are better for the survival of more delicate specimens. Rawlins (1992: 56) suggests that glycerol fades many stains. Ramanna (1973: 103) found that Euparal was suitable for preserving fluorescence of Aniline Blue in plant material such as pollen tubes. Partial bleaching with ammonia and hydrogen peroxide may be required if the specimens are intransigently opaque due to dark sclerotisation as in some Aleyrodidae homopteran bugs.

2.10 Microscope slide mounting methods.

To select the most suitable slide mountant one must refer to literature covering the study of the organisms concerned (as listed in the bibliography) Great care is needed in selecting methods to balance the requirements for visibility with the longevity of the mount and the specimen within the mount. For example, doubts about the permanence of gum chloral mounts have now been widely published (Noyes, 1989, Upton, 1993 and see discussion below).

The mountant used in a slide preparation should always be noted on the slide label. Names used should also indicate a specific recipe if possible, and the name of the preparator and the date. Gum chloral mounts have indiscriminately been called Berlese, Hoyer's or Faure's even when the recipe is quite different from the original in the concentration of constituent parts. A deteriorating slide without exact details of the mountant can prove difficult to repair, and may involve trial and error soakings in different solvents which may damage the specimens and waste much time. Often slides were made as temporary mounts and were not meant to last, but a change in status of the specimens, for example to 'type' series, dictates the use of a more permanent mount so as to preserve the type for future taxonomic study.

Many slide mountants require a period in a curing oven at 30-40 degrees centigrade to harden the mount by evaporation

of the solvent especially if the slides are to be stored vertically as a soft mountant will creep under the influence of gravity. If the mountant is soft, later handling can also move the coverslip possibly rolling and ruining the specimens. The use of wire springs to hold the coverslip, as described by Wagstaffe & Fidler (1970: 197), is employed by some workers to compress the specimens while the mountant hardens.

2.11 Ringing the coverslip.

Liquid, glycerine and gum chloral mounts need to be ringed round the edge of the cover slip to seal the mountant and prevent its escape, loss through evaporation, or oxidation through contact with air. Excess mountant from round the edge of the coverslip should be removed by scalpel before ringing. Ringing of round mounts is easily accomplished using a mounting turntable (Fig. 11). The slide is clipped onto the revolving stage and a small amount of ringing medium such as Euparal, Glyceel, Glyptal (Travis, 1968) or Murrayite is applied with a fine brush as the slide revolves. Wu (1986) describes his method of applying Glyptal to nematode 'Hoyer's' mounts using a plastic bottle. The many sealants used are listed in Appendix 1. Cutex and other brands of nail varnish have been used for both ringing and as a mountant. Wells (1978) describes nail varnish used with butyl acetate-acetone to mount mollusc radulae and pollen grains. The use of more than one coverslip on a slide (Figs. 8 & 9) can make ringing difficult and tedious.

If the mountant is prone to shrinkage, ringing may not stop the ingress of air as any stress applied to the ring or the coverslip may cause them to fail. A further application of the mountant at a later date may be necessary to replace the shrinkage loss, before a ring is applied. Resin mounts do not normally need ringing as the solvent in the resin needs to evaporate in order for the mountant to harden.

2.12 Labels.

Bridson & Forman (1992) state that gummed labels for microscope slides should be of "archival" quality, be they foil backed, gummed self adhesive or manually glued. Non-archival gummed and self-adhesive labels have often been used in the past and have sometimes become detached or become transparent after a few years. Where slides are stored vertically in contact with each other, the most suitable labels consist of thick card squares glued to the glass with a polyvinyl acetate glue. These have the advantage that they also act as 'spacers' between the slides to protect the coverslip and the slides can be stacked for drying without fear of them sticking together and becoming useless. A diamond-tipped engraving scribe can be used to write the locality details and accession number directly on the glass slide beneath the labels. This provides the means of maintaining the identity of the slide when labels become accidentally detached or the writing on labels fades. The Microlepidoptera genitalia slides, plant chromosomes and pollen slides and Protozoa slides (in the NHM collections) are marked in this way.

2.13 Immersion oils.

For viewing microscope slides at high magnification, an immersion medium is required between the microscope's high power objective lens and the coverslip. Such immersion oils are available from a number of microscope manufacturers and usually have a refractive index close to glass at about 1.5. Immersion oil should be cleaned off the slide after examination as it forms an unsightly smear on the glass and might cause deterioration of the mountant or the sealing ring (M.G.C., 1992: 52).

2.14 Storage.

Collection storage has often been dictated by that already in use for major donated collections, so that, for example slide storage at the NHM is not standardised. Storage of slides can be horizontal (slides laid flat) on shallow slats and trays in boxes (Fig. 12) or cabinets; or vertical within slotted boxes (Fig. 13) or drawers (Fig. 14) which are a little deeper than the width of the slides. At the NHM slides are stored in a variety of wooden and metal cabinets, either vertically orientated in drawers (about 800,000 slides, mostly in Entomology Department) or horizontally in wooden (Fig. 15) and metal cabinets (Fig. 16) and in boxes with slats (about 1,100,000 slides).

When considering the particular needs of a large microscope slide collection one can quote Mound (1992: 10) who describes the 400,000 NHM Homoptera and Thysanoptera slides combined, as a solid block of glass of 12 cubic metres. This weighs about 8 metric tonnes and exerts a load of 0.3 tonnes/square metre or about 4 kiloneutons/square metre. Such weight must be carefully considered when floor loadings are concerned! The regulation minimum floor loading is 5 kiloneutons but the NHM Entomology block floor has been recorded as capable of carrying 17 kiloneutons (Colin Farniloe, pers. comm.).

One must decide on the right type of cabinet in which to store slides. In the NHM, generally the thinner the slide mounts, the more likely they will be stored vertically so that the slide collection can become a self indexing system. The thicker the mounts, the more likely they will "creep" under the influence of gravity when stored vertically, so such slides are better stored horizontally. Liquid mounts are also stored horizontally as the seal is more prone to damage by jostling with other slides and the specimens will sink to the lower edge and be damaged against each other or against the edge of the mount. However, slides stored horizontally take up more space.

In the NHM, standard wooden cabinets (so called Hill units) with horizontal slide drawers hold 5250 slides (Fig. 15) and those with vertical slide drawers can hold 10,000 slides when full (Fig. 14). Before storing slides vertically they must be baked hard in an oven at 30-40 degrees centigrade to avoid the mountant "creeping". The vertically stored Homoptera slides (Entomology Dept.) are carded (the thickness of the card label on each side of the mount gives some protection) and enveloped to give extra protection from dust and physical damage (Fig. 18). Previously, manilla envelopes specially manufactured were used to house slides individually (Fig. 18), though the time spent duplicating information was considered excessive, but is still used for

type and other important specimens especially when that material is loaned, so that the information remains within the collection. In the past, plastic sleeves have been manufactured from unplasticised polyvinyl chloride which is not of archival quality and which are turning yellow and becoming brittle. New plastic sleeves are being made of archival quality polyester (by Preservation Equipment Ltd.) Such vertically stored collections can be arranged taxonomically with closely related families, genera and species together (and with associated indices), or they can be arranged alphabetically (by family, genus and species in unrelated order), and because they are thin and card-like, they can be their own index obviating the need for a separate card index (Fig. 16). With vertical slide storage it is possible to incorporate short bottles of specimens in spirit, pinned dry specimens in unit trays and dried host plant samples in the same drawer, as in the NHM Diptera collection (Fig. 19). Unit trays holding about 25 slides each are used in some vertically stored collections to facilitate easy removal of slides (Fig. 16). At the Royal Botanic Gardens at Kew, plant anatomy slides are housed in Gallenkamp metal clips (4 to a clip) which fit vertically in standard card file drawers, the slides standing upright on end.

As with most natural history collections, microscope slides require a controlled environment. There are four lines of defence — the sealant ring, the envelope, the cabinet and the room in which the cabinet is housed. The Museums & Galleries Commission's Standards in the care of Biological Collections (1992: 52) makes little mention of the specific needs of microscope slides, except to list them as being prone to damage from variation in relative humidity. In high relative humidity, fungal attack can occur in dry, liquid and other aqueous mounts and creep and sweating can occur in aqueous mounts. In low relative humidity, discolouration and cracking can occur in these same mountants (see section on Gum chloral mounts below.) Sometimes, slides can be damaged when they are lent to institutions in which the environment is not controlled or when they are subjected to varying conditions in transit. Slides should be stored in total darkness except when being examined as high light level is suspected in the deterioration of gum chloral mounts and in the fading of stains. Routine freezing of incoming parcels to kill insect pests may damage aqueous micro-slide mounts.

3. Which microscope slide mountants should be used ?

The general museum literature gives little in the way of guidance on the merits of different microscope slide mountants, with no mention of drawbacks in Walker and Crosby (1988) or in Stansfield (1992: 443) for example. General works covering the choices in the making of microscope slide mounts for biological specimens include Gray's *Microtomist's Formulary and Guide* (1954), Wagstaffe & Fidler's *Preservation of Natural History Specimens* (1970) and Knudsen's *Collecting and Preserving plants and animals* (1972). Often, to find the latest preferred method, one has to consult the texts for the specific organisms with which one is dealing. Some preparators have no knowledge of why they used their chosen mountant, apart from the fact that they had always used it by tradition, which might indicate that the mountant is suitable but possibly not the best available. Individuals working in relative isolation

may not be aware of better techniques which are worth experimenting with. Recipes and techniques have been published in 'special interest' journals which are not read by other workers studying in different disciplines. Specialists also have a tendency to develop their own, often complex techniques, with little regard to archival quality, compatibility, standardisation, or ease of future handling, curation or conservation.

Rawlins (1992: 56) suggests that there are two categories of mountants, the permanent and the semi-permanent (which do not set hard). Gutierrez (1985: 352) states that "no mounting media are fully satisfactory" for spider mites. Lillie et al, (1953: 71) came to a similar conclusion after carrying out an exhaustive survey of histology mounting media. Many workers would agree with this statement but most workers who wish their slide mounts to remain "permanent", choose a favourite mountant which both suits their viewing requirements and preserves the specimens for future research. Such workers have frequently made up their own mountants and given them "pet" names, often not indicating the recipe used for a particular slide and sometimes not bothering to publish the recipe. There is much confusion as to the correct recipes for mountants and when they were first used, as discussed by Upton (1993) for gum chloral mounts. Some proprietary brands of mountants are and have been made to secret recipes whose names or recipes have been changed and have been copied by others, which has added to the confusion. Since the last century, collections have grown and have been amalgamated into major museum collections, some of which now require conservation. The conservator may be faced with the problem of not knowing what the deteriorating mountant consists of, even if the name of the mountant is written on the slide, as it may not be the published recipe. With the passage of time, the evidence of degradation of some mountants has slowly and, sometimes too quickly, become apparent, adding another dimension to the discussion about the best refractive indices and the short term effect of the mountant on the objects.

3.1 Refractive indices (see Appendix 2).

"The ideal mounting medium for stained preparations should have the same refractive index as the mounted object." (Pantin: 1964, 21). Conversely, if the specimens are colourless, as are diatoms or mites, then visibility in bright field microscopy is enhanced by a difference in refractive index. A diatom of approximate refractive index of 1.4 when mounted in a mountant of 1.5 will have a visibility proportional to the difference between 1.4 and 1.5 (Fleming, 1943: 34). Knudsen (1966: 500) states that diatoms and mites become 'lost' in Canada balsam because the refractive index is close to that of the mountant. Acarologists such as Norton of Syracuse New York State University, continue to use gum chloral mounts because they offer a "better" contrast with a lower RI. Diatom researchers use mountants with a higher RI of 1.6-1.7 (such as Naphrax) than diatom silica at 1.43, (McLaughlin 1986: 287). The use of stains and of phase contrast has largely overcome this problem. Walker & Crosby (1988: 22) mention that Canada balsam is "particularly recommended as a mountant because its refractive index (1.53) is very close to that of glass". They

do not refer to the problem of reduced visibility of small and transparent organisms in bright field. It is important to remember that refractive index changes with temperature of the mountant and with the percentage loss of solvent from it. Canada balsam dissolved in xylene is 1.497 and dries to 1.532 (Loveland) and so the optical quality of the mountant changes from when the slide is first made to later inspection.

3.2 Natural resinous media.

3.2.1 Canada balsam (RI = 1.52-1.54).

Canada balsam was first described as a suitable mounting medium for the new science of transmitted light microscopy by Andrew Pritchard in the 1830s. It is the most widely used mountant because of its proven archival quality, with a track record of over 150 years, and does not crystallise or absorb moisture as do gum chloral mounts. Eastop (1984: 248) observed that Canada balsam slides of aphids made by Francis Walker in 1847 show no signs of deterioration except for yellowing (Fig. 20). This yellowing is demonstrated in Fig. 21 where one yellow Canada balsam slide is compared with 3 colourless Euparal slides of the same age (20 years). Mound & Pitkin (1972: 122) state that Canada balsam is the only mountant known not to deteriorate when kept for many years in a variety of climates. Noyes (1982: 329) states several million years as the longevity of Canada balsam, comparing it with natural fossilised amber. Because Canada balsam is similar to amber, it has been used to mount insects in amber on to microscope slides (Grimaldi, 1993, 46). Figure 22 is of the infamous 'fossil' latrine fly hoax discovered by Dr Andrew Ross and reported in the *New Scientist* (Palmer, 1993: 4) where the fly was mounted in Canada balsam within a manufactured pocket in a genuine piece of Baltic amber. Canada balsam can be acidic, which will erode calcium carbonate and fine structures on some organisms and minerals, so one should be certain to select a neutral balsam as described by the manufacturer for slide preparations of coccoliths and ostracods.

Hood (1940), who mounted over 50,000 microscope slides, came to the conclusion that Canada balsam was the only mountant to use. Similarly Cooper (1988: 228), after many years experience in the NHM Histology section, used only Canada balsam. Saito & Osakabe (1992: 427) considered Hoyer's to be only temporary and published (1993: 593) an improved method for mites using Canada balsam, fixing in methanol-acetic acid and with x-terpineol, but they do not mention the refractive index. Figure 7 shows a deep mount of a whole Trichopteran in Canada balsam which has darkened with age and has thus reduced the visibility of the specimen. Specimens as large as this are better preserved dry on pins or in spirit. Yellowing and darkening was not considered a serious problem by those commenting on balsam in the NHM survey. The tedious rinsing through three or four, more concentrated, alcohols to dehydrate specimens has been replaced by a single rinse in glacial acetic acid so that the Canada balsam technique is now less time consuming. Rawlins (1992: 56) relegated Canada balsam as having been widely used at one time as it is strongly "autofluorescent". He also points out the harmful solvents which constitute a health hazard such as xylene, might limit the choice of mountants in UK museums which

must conform with Control of Substances Hazardous to Health (COSHH) regulations. The use of non toxic solvents for Canada balsam instead of xylene (possibly Cellosolve or HistoClear) would alleviate the safety problem but might cause other problems such as slower hardening rates and premature darkening. Some workers have reported deterioration in Canada balsam such as "crazing" by Green (1995: 162) which might be due to incorrect preparation of specimens or a genuine problem in a minority of balsam mounts.

3.2.2 Phenol balsam.

Advocated by dipterists working on small flies, this variant of the Canada balsam technique, replaces xylene with phenol as clearing agent and thinners (Wirth & Marston, 1968). It is more convenient and "produces the best slides". NHM dipterists have considered using this as the preferred mountant for Ceratopogonidae and, possibly, the Simuliidae. Phenol can be used as relaxant, clearing agent and dehydrating agent in one soak and was considered less dangerous than xylene. The author's discovery of black Canada balsam slides in the aphid collection (Fig. 23) may indicate future problems with the use of phenol balsam. Specimens cleared in chloral phenol and then mounted in Canada balsam have turned black with similar cuticular disruption as that found in gum chloral mounts, unlike the contemporary material cleared with KOH in balsam, so this suggests a link with phenol. These slides blacken from the edge inwards which may suggest atmospheric oxygen being involved so ringing may be required to prevent this from happening. Drs Richard Lane and John Boorman (pers. comm.) have suggested the wider use of phenol balsam as a mountant in the NHM Department of Entomology slide collections but this may be risky in the light of the author's experience.

3.2.3 Euparal (RI = 1.48).

The Euparal recipe is now a trade secret but early papers list the possible ingredients as being eucalyptus oil, methyl salicylate, camphor, sandarac, and possibly paraldehyde. Euparal is manufactured in Germany and Britain (ASCO Laboratory, Manchester) and the two products have been found to differ. It has been reported that workers at the Smithsonian Institution have found that the German product does not match the standard of the ASCO Euparal.

Rawlins (1992: 56) lists Euparal as being more popular as a permanent mountant than Canada balsam. Euparal is widely used and in the NHM its use is second only to Canada balsam, being used for cytology mounts, plant mounts (also at The Royal Botanic Gardens at Kew) and for some insects. The virtues of Euparal were extolled by Imms (1929: 166) as not yellowing with age like Canada balsam and as having a lower refractive index of 1.48. Thus structures which are too transparent in Canada balsam can be seen in Euparal with bright field microscopy (Fig 24). This is the main argument used by lepidopterists at the NHM (see discussion on Canada balsam for comparisons). Euparal is regarded as being a good permanent preservative, proven over the passage of time (for example, over 30 years in the NHM Microlepidoptera collection), of consistent quality, safe, quick and easy to use, good optically with low

refractive index and drying quickly. In the NHM many slide preparators (as reported in the mountant survey part of the MA project), commented about Euparal being better than Canada balsam, "not requiring the use of carcinogenic xylene". Euparal seems to be the most popular alternative to balsam, especially if COSHH regulations preclude the use of Canada balsam in either xylene or phenol mixtures. Hood (1940) stated that Euparal was unsuitable because it quickly developed a meniscus which can damage fine structures when specimens were being arranged in the mountant, bubbles took longer to clear from the mount if at all, and that Euparal slides were prone to crystallisation although in the author's experience this does not occur. This might be due to incorrect preparation technique or possibly might involve the German product. Carolyn Lowry (pers. comm.) has experimented with Euparal and found that some fine structures of Simuliid flies become too fragile in the slide making process to make good mounts. Theresa Howard has found that using Cellosolve instead of Euparal Essence keeps such fine structures supple so that they can be arranged more easily. At the RBG at Kew, HistoClear is used instead of Cellosolve with the same benefits.

3.3 Synthetic resinous media (plastics).

These are rarely used at the NHM due to the known shrinkage and crazing in plastic mountants such as DePeX and polyvinyl lactophenol.

3.3.1 Cellofas = Carboxy Methyl Cellulose (CMC) (RI = 1.428).

Used in the NHM Palaeontology Department for mounting fossil specimens. Stehr (1987: 16) questions whether CMCP-9 is suitable as a permanent mountant. Knudsen, (1966: 501) describes how Turtox CMC can be used to kill, clear, stain (CMC-S) and permanently mounts specimens from water or alcohol in one process. Hobson & Barse (1981: 6) refer to its use for polychaetes and mention that it contracts when hardening so that extra mountant must be added later before the slide is ringed. Clark & Morishita (1950: 789) used CMC for mites because of its low refractive index and because it did not crystallise.

3.3.2 DePeX (Distrene plasticiser xylene) (RI = 1.53).

Rawlins (1992: 56) lists DePeX incorrectly as being a permanent mountant. Gurr (1956: 42) mentioned that DePeX suffers from "a considerable degree of shrinkage when drying" and suggests that it should be applied liberally to the slide to allow for this. What once seemed a perfect mountant with suitable refractive index has now proved not to be permanent because of continued shrinkage causing disruption of specimens and loss of coverslip with a change in refractive index and severe optical distortion within specimens. Only one collection of (aphid) chromosomes is still mounted in DePeX and this is of secondary importance to the collection of photomicrographs.

3.3.3 Dimethyl hydantoin formaldehyde (DHFM) (RI = 1.45-1.46).

This synthetic resin which is miscible with water has recently been suggested as a suitable mountant by Dr M.

Alonso-Zarazaga of the Museo Nacional de Ciencias Naturales, Madrid (pers. comm.). Trials may be carried out with mites as an alternative to Hoyer's mountant. Nematodes (Smith, 1966: 177), beetle genitalia (Cooter, 1991: 57) and turbellaria (Steedman, 1958: 451) have been mounted in DHFM. It is reported not to shrink but otherwise there is no record of longevity.

3.3.4 Naphrax (RI = 1.71).

Of the naphthalene resins only the toluene-based Naphrax is still used in the NHM for mounting diatoms which need a large contrast in refractive index to make the surface detail of the diatom's silica (RI = 1.43) visible (Green, 1995: 163). McLaughlin (1986: 285) discusses the requirements for a good diatom mountant. Ragge (1955: 8) found that the tracheae, supported by chitinous taenidia, of orthopteran wings were almost invisible in Canada balsam, and consequently he changed to using Naphrax. After 50 years some darkening is evident (Fig. 25).

3.3.5 Permout (RI = 1.51).

Used for mounting Porifera in the NHM and recommended together with Numount, for their clarity and non-shrinkability. Dr Eric Metzler (pers. comm.) reports that his twenty year old lepidoptera genitalia mounts in Permout have crystallised but that they can be recovered by soaking in xylene. Otherwise there is little in the literature reporting on the permanence of Permout.

3.3.6 Histomount (RI = 1.49-1.50)

Although this mountant has been little used at the NHM, Donald Quicke has seen no deterioration over 10 years and the importers National Diagnostics report no cases of deterioration in 20 years, unlike with the less successful Omnimount. Histomount is at present being tested by the author. This mountant can be thinned with the unrelated Histoclear which is considered a safe solvent being based on orange oil.

3.3.7 Polyvinyl Lactophenol (PVL) (RI = 1.43-1.44).

Huys & Boxshall (1991: 451) suggest that polyvinyl lactophenol "is not suitable for type collections because specimens (of copepods) are slowly over-cleared in the mount, the mountant is slowly replaced with rosettes of long thin crystals and it often dries out if not sealed". The NHM Zoology department uses PVL for mounting copepods and reptile scales but otherwise it has fallen out of favour, although the Central Science Laboratory of the Ministry of Agriculture still uses the Heinze formula (Heinze, 1951: 177) for mounting thrips and mites. Bink (1979: 160) mentions that some 18 year old polyvinyl lactophenol preps were in good condition but others were crystallising and turbid. Increasingly, polyvinyl lactophenol is being considered as semi-permanent at best and thus not suitable for important material as it continues to shrink (Fig. 26).

3.4 Gum chloral mountants (RI = 1.48).

In the NHM survey, "Berlese" and "Hoyer's" gum chloral mounts were stated as being the best optically for the groups concerned as well as traditionally used. This group of

mountants consists of many closely related recipes including gum Arabic, phenol, glycerol, lactic acid, sugars and salts. Upton (1993) discusses these recipes in detail and discusses how many differing recipes have been confused and authors misquoted. Some describe "Berlese" as being only semi-permanent and advise the use of Canada balsam (Hodkinson & White, 1979: 3) and Euparal (Freeman, 1987: 196). Likewise, Kozarzhevskaya (1968: 147) lists "Berlese" only as a temporary mount for scale insects. Stehr (1987: 16) questions the permanence of "Hoyer's" and states that, under humid conditions, unringed "Hoyer's" slides will loosen and slip. Noyes (1982: 329) states that resinous mountants (such as Canada balsam) withstand a wider range of climatic conditions than water soluble "Hoyer's" and "Berlese" which are not only sometimes unstable but frequently cause distortion to such insects as thrips. Bink (1979: 160) mentions "Faure's" as becoming sticky in humid climates.

3.4.1 Crystallisation.

Gum chloral mountants deteriorate by dehydration of the mountant with the resulting formation of small white and opaque chloral hydrate crystals (Fig. 27). These almost always form from the outer edge and advance towards the centre, overwhelming and obscuring the specimens. This occurs when the slides have not been ringed to avoid dehydration. Specimens can be rescued from a crystallised slide as the cuticle is not chemically eroded, although there can be some physical damage from the growing crystals. Crystallised slides can sometimes be saved by rehydrating the gum chloral mountant in a warm, moist atmosphere with Thymol to avoid fungal infection. Otherwise, specimens can be soaked out of the gum chloral in water, dehydrated in glacial acetic acid and remounted into Canada balsam (Martin, 1987). Disney (1988: 106) suggests an excess of glucose of over 8% of the mountant is the cause, but Freeman, (1987: 196) quotes Keith Harris who blames the poor quality of modern gum Arabic. It is more likely due to a higher percentage of chloral hydrate in the recipe such as in Imms' "Berlese". If glycerine is used in the recipe instead of glacial acetic acid and glucose syrup is present, a granular opalescence can develop in the mountant which obscures fine hairs and cuticular details (Eastop & van Emden, 1972: 8).

3.4.2 Blackening.

Blackening was not listed as a problem with gum chloral mountants by Upton (1993) but this has become the most serious problem within the NHM Entomology department slide collections and can be irreversible (Figs. 8 & 28). Within the Aphidoidea collection, the first sign of deterioration is a pinkish tinge emanating from the specimens, which rapidly turns bluish and then opaque black across the whole mountant, within a period of as little as six months. The cuticle of the aphid becomes transparent and then disappears, sometimes before the slide has become opaque. The cause of this blackening has been attributed to inadequate rinsing of the macerating and clearing agents, potassium hydroxide and chloral phenol respectively, from the specimens. The presence of light and high temperature have also been suggested as contributing factors. Other mounts made in the 1970's which were cleared with chloral

phenol instead of KOH have distorted due to osmotic differences between mountant and body contents. Stuart Harbron (pers. comm.) suggests that the blackening may be a polymerative oxidation of phenol, started by a single molecule of an oxidant such as KOH and possibly also triggered by light. Blackening starts within the specimens where phenol has been entrapped when the specimens were mounted from chloral phenol to the gum chloral mountant. Womersley (1943: 181) uses phenol in the gum chloral mountant to try to prevent crystallisation but the presence of 25% phenol may be a sure invitation for blackening to occur. At the NHM, expressly because of blackening, Hemiptera are now mounted in Canada balsam after clearing in KOH with careful rinsing in 30% ethanol and dehydration using glacial acetic acid with another careful rinsing in clove oil as standard procedure. The problem of Canada balsam having the same refractive index as insect cuticle is overcome by the use of phase contrast microscopy instead of bright field. Specimens which have not been cleared in chloral phenol before mounting in Gum chloral may not turn black but this possibility has not been fully investigated.

Other gum chloral slides turn brown at a much slower rate with a slower rate of erosion of the insect cuticle which may be related to the above. Dr Victor Eastop has suggested (pers. comm.) that these specimens were initially cleared with lactic acid and not potassium hydroxide so that the chemical reaction may be a different one. A different form of blackening occurs in some of the Stroyan aphid slides where the mountant darkens from the Murrayite ringing medium. This blackening does not damage the specimens.

The Lewis collection of phlebotomine and simuliid dipteran flies consists of 20,000 "Berlese" slides, which are also turning black. This blackening (Fig. 8) has been attributed to a chemical reaction with the Euparal ringing mountant and has been noted as starting at the edge and working inwards (Lane, 1993: 112). Disney (1988: 106) suggests the blackening is attributed to residual preservatives such as phenolics and formaldehyde within specimens. This mountant dissolves in acetone and the specimens can be remounted successfully. There is no evidence of cuticular disruption, even in the most blackened slides. Lewis quotes the Puri recipe in Kirk & Lewis (1951: 500), which could be the problem mountant he used for his collection, and later in Smith (1973: 173) he also quotes the Stroyan (1949) recipe which he might have found to be better. With aphid Berlese slides, even if the blackened Berlese can be restored, the insect cuticle is still irrevocably damaged and a programme of remounting material not yet damaged is much better than restoring already blackened damaged specimens.

After the long discussion in print about the effectiveness of Canada balsam (Noyes) versus the permanence of "Berlese" (Disney), Disney (1994: 387) finally concedes, quoting Upton (1993) and admits that there are problems with the permanence of gum chloral mountants but advises the use of Euparal for a majority of mounts and a few in "Berlese" to illustrate the fine characters such as the Dufour's crop mechanism of many phorid fly females.

Disney (1989: 48) risks the loss of permanence for bright field, visual access to museum specimens. His comment that remounting type specimens every 50-100 years is a small

price to pay, might more likely be every 15-30 years and the cost of employing a trained worker to undertake the job could be much reduced by using Canada balsam and purchasing a phase contrast microscope! Each time fragile specimens are remounted, there is a high risk of inadvertent or other damage taking place. Some workers have used gum chloral mountants for many years and have not seen any such problems possibly because they do not use phenol as a clearing agent (Zhang, pers. comm.). However, in the face of the well known problems, only mites and some small dipteran groups are still mounted in gum chloral mountants at the NHM.

3.5 Liquid mounts. (Fig. 17)

Some workers consider that liquid mounts can be permanent if ringed with a permanent ringing medium. But ringing media can shrink as they age and behave differently than when used as mounting media under a coverslip. A thin ring can crack and allow the liquid to evaporate. Gutierrez (1985: 352) considers liquid mounts as only temporary. Dieguez & Montero (1992: 315) found that 90% of the diatom liquid formaldehyde slides of E. Caballero Bellido made between 1891 and 1927, had survived in good condition. Formalin mounts are no longer used at the NHM due to risk of drying out (25% of formalin mounts have been lost in the Rotifera collection and no staff are available for conservation work) but Ealing Hospital uses formalin sealed in paraffin wax for permanent collection of tissue biopsies (J.D. Arnold pers comm.).

3.5.1 Glycerine-glycerol (RI = 1.40-1.47).

Glycerine and glycerol variants are the most common liquid mounts at the NHM, being used to preserve pollen and spores in the Botany Department, Nematoda, Rotifera and Bryozoa in the Zoology Department and miospores and dinoflagellates in the Palaeontology Department. It is used in liquid mounts either on slides or in glass or gelatine capsules where the three dimensional nature of the structure must be fully preserved. Spence (1940: 144) reports that it prevents mould growth and does not dissolve calcium carbonate. Glycerol readily shrinks and expands with variation in relative humidity which could stress coverslip and specimens unless totally sealed. As with other water soluble mountants it was chosen because it was cheap, safe, quick and easy to use with little preparation needed for the specimens, and it is inert and good optically. Wiles (in manuscript) discusses his technique for double coverslip gelatine mounts using aluminium Cobb slides for water mites.

3.5.2 Karo corn syrup (RI = 1.47).

This is used by the NHM Botany Department to mount plankton and parts of larger marine and freshwater algae. Karo corn syrup or dextrose is commonly available and the technique is described by Monk (1938: 174) and Knudsen (1972: 23). Taft (1978: 263) uses glucose in a similar technique. It is chosen because it is cheap, safe, quick and easy to use with little preparation needed for the specimens, "algae can be placed straight into Karo syrup from sea water", and it is inert and good optically. An effective seal is required to stop water loss from the mount.

3.5.3 Lactophenol.

Huys and Boxshall (1991: 451) recommend lactophenol as the best mountant for Copepods as long as the slides are effectively sealed. In the NHM both Copepods and Crustacea are mounted in lactophenol.

3.6 Ringing media.

The resin mountants do not need ringing, although White (1992: 29) states that "it is not advisable to store Canada balsam slides vertically unless they have been ringed with Glyceel or a similar ringing compound". Gray (1954) suggests that Canada balsam be ringed to stop it darkening by atmospheric oxidation. Disney (1988: 106) suggested that Euparal, when used to ring "Berlese" mounts, causes discolouration of "Berlese" over time. Within the Stroyan aphid collection, discolouration of the gum chloral mountant is probably due to direct contact with Murrayite and not Euparal. Disney advocates Glyceel, Trycolac and "ladies' nail varnish with or without colour". Paraffin wax is a secondary sealant which can be smeared over the first sealant to reinforce the seal. Canada balsam, red Glyptal insulating varnish and Zut have, with other shellac and asphalt based cements, been used successfully at the NHM.

3.7 Adhesives.

The NHM Palaeontology and Mineralogy laboratories use a number of synthetic adhesives to glue sections and small specimens on to microscope slides. This specialist application is unrelated to the main requirements of natural history specimen mounting. Refractive index and permanency are important as also are the pH and the chemical constituents of the glue or mountant as such might erode or even destroy the specimen. The following media are presently used; Araldite epoxy resin, Cellofas, Crystalbond thermoplastic cement, Dymax ultra violet bonding resin, Elvacite, Epotek epoxy resin, Lakeside, Loctite ultra violet bonding resin and Petropoxy epoxy resin. Apart from Crystalbond which is a temporary glue, there has been no criticism in the survey regarding deterioration, and only the passage of time will show which are the most suitable.

4. Literature survey & Bibliography.

A literature search was undertaken prior to the survey. Relevant papers on slide mountants in both microscopy and taxonomic journals were collected (and later augmented by the references given in the returned questionnaires). Many taxonomic papers had notes on mountants used for the organisms but such details were not reflected in the titles of the papers and so were not accessible to the computer searches which were carried out. A total of 250 books and papers were studied and included in the lengthy bibliography (of which only 28 were volunteered through the questionnaire). This bibliography and the list of 150 mountants and adhesives and published recipes gleaned from the papers (Appendix 1) are as important as the survey itself as few such lists exist. Gray (1954), Lillie et al. (1952) Barbosa (1974), Loveland & Centifanto (1986), and Upton (1993) study mountants in detail, Gray listing 243 mountants and adhesives of which many are archaic. Comprehensive works reviewing relevant mountants and slide mounting

techniques required for the study of specific, or for many different groups of organism included Hood, 1940 (insects), Johansen 1940, (plants), Wagstaffe & Fidler, 1970 (invertebrates), Knudsen, 1972 (plants and animals), Martin, 1977 (insects), Smithers, 1981 (insects), Steyskal, Murphy & Hoover, 1986 (insects) and Higgins & Thiel, 1988 (marine invertebrates). With the published information the author could then compare information from the returning questionnaires and draw conclusions from the collected data. Within this bibliography almost every organism is mentioned and the aim was for the reader of the dissertation to be able to choose a mountant and preparation technique suitable for their requirement.

5. Conclusion. Which are the best mountants and what future have slide collections?

How safe will the microscope slide collections be in the future? Legal obligations protect collections from disposal but what of the insidiously deteriorating mountants? The preparator must carefully consider all the risks involved and balance the current needs for optical quality with long term preservation.

With little guidance in the general museum literature on the merits of different mounting media, one must refer to the specialised literature covering the study of specific organisms (see the bibliography) to decide on the most suitable mountant. Descriptions of latest techniques are often difficult to find, as they are often appended to papers on taxonomy in specialist journals which are not widely available. Organisms react in different ways to different mounting methods. Some become too fragile or too rigid in the process to make good mounts.

Discussion will continue about the optical requirements of different organisms but the consensus of opinion is turning away from experimentation with new mounting media (except in the high powered experimental end of electron microscopy) as more and more slides are found to be deteriorating. It must be stressed that many mountants should not be considered permanent and that routine monitoring of collections should be standard procedure as with spirit and dry collections. Likewise, environmental conditions should be constant with regular monitoring as light, desiccation and freezing can have deleterious effects on many mountants. Other synthetic resin and gum chloral recipes which produce perfect slides both at present and in 20 years time, may deteriorate in 50 years time. The cost of employing staff to remount slides is expensive and may not be considered affordable now or in the future. This may also apply to the routine re-ringing of liquid or jelly mounts but some organisms demand the use of less permanent mounts, such as glycerol and karo syrup. The newer plastic adhesives used for plant and mineral sections are as yet untested by the passage of time and require staff to monitor them. The NHM has stopped the practice of clearing specimens in chloral phenol before mounting in gum chloral mountants and will continue and increase the use of Euparal and Canada balsam, especially if the latter can be thinned with the safe solvent Histoclear instead of xylene as they need relatively little future conservation monitoring. These two mountants should be used for primary types especially with the use of staining of specimens and phase contrast microscopy to overcome the

refractive index problem. Field tests of new mountants will continue with Histomount, dimethyl hydantoin formaldehyde and Entellan showing signs of being suitable preserving media.

6. Acknowledgements.

This paper is a revised version of part of a dissertation accepted for the degree of MA in Museum Studies at the Institute of Archaeology, University College London, 1994. I thank Elizabeth Pye, head of the Conservation and Museums Studies Department (UCL) for permission to publish, and Geoff Stansfield for supervising the project. I also wish to thank the following friends and colleagues for their help at the NHM; Zoe Adams, Dr Anne Baker, Dr Peter Barnard, Dr Roger Blackman, Dr John Boorman, Emma DeBoise, Dr Victor Eastop, Theresa Howard, Dr Richard Lane, Carolyn Lowry, Jon Martin, Dr Lawrence Mound, Dr John Noyes, Dr Brian Pitkin and Kevin Tuck of the Entomology Department; Karen Webb, Alison Paul, Len Ellis and Jenny Moore of the Botany Department; Dr Geoff Boxshall, Oliver Crimmen, Sheila Halsey, Mary Spencer Jones & Dr Alan Warren of the Zoology Department; William Lindsay & Dr Andrew Ross of the Palaeontology Department; Chris Jones of the Mineralogy Department; all others not mentioned above for filling in the questionnaire; Dr Chris Malumphy at the Ministry of Agriculture, Central Science Laboratory, York; Dr P. R. Wiles at the School of Sciences of the University of Buckingham; Peter Gasson and Tim Lawrence of the Jodrell Laboratory at the Royal Botanical Gardens, Kew; Dr Sebastiano Barbagallo at Catania University, Sicily; Dr Roy A. Norton at the University of New York at Syracuse, U.S.A.; Simon Moore of Hampshire Museums Service; Dr Stuart Harbron of the UCL Chemistry Department for helpful discussions on possible reasons for gum chloral mountant deterioration; Julie Harvey and Carol Gokce of the NHM Library for much help in chasing references and Geoffrey Kibby of the International Institute of Entomology for computer file conversion.

7. Appendix 1: Mountant recipes

When faced with a set of deteriorating microscope slides one needs to find out which mountant was used and what active ingredients and environmental factors have caused the deterioration. Here follows a list of mountants with recipes when known. Some proprietary brands have "secret" recipes and are chemically unidentified. Recipes are taken from the literature and the references are cited (within brackets). Mounting media which are underlined are presently in use at the NHM. Sources of supply are given [in square brackets].

NATURAL RESINS

Canada balsam = resin of *Abies balsamea* in Xylene or native.

(Barbosa, 1974: 86) (Baylak, 1986: 311) (Clastrier, 1984: 175) (Cooper, 1988: 228) (Cutler, 1978: 6) (Essig, 1948: 16) (Gates Clarke, 1941: 150) (Green, 1995: 161) (Hangey, 1985: 147) (Heming, 1969: 323) (Hood, 1940: 53) (Huys & Boxshall, 1991: 451) (Johansen, 1940: 110) (Knudsen, 1966: 501) (Kozarhevskaya, 1968: 146) (Martin, 1983: 411) (Moore, 1979: 494) (Mosely, 1943: 231) (Mound & Pitkin,

1972: 122) (Noyes, 1982: 329; 1989: 2) (Oldroyd, 1970: 137) (Palma, 1978: 432) (Purvis, 1964: 106) (Rattanaarithikul, 1982: 148) (Rawlins, 1992: 56) (Remaudiere, 1992: 185) (Richards, 1964: 963) (Saito, et al., 1993: 593) (Smithers, 1981: 77) (Stehr, 1987: 16) (Wagstaffe, 1970: 174) (Wilkey, 1990: 349). [BDH (Gurr) and Hopkins & Williams].

Phenol balsam = Canada balsam thinned with phenol alcohol (without Xylene). (Barbosa, 1974: 86) (Boorman & Rowland, 1988: 65) (Lane & Crosskey, 1993: 113) (Wirth & Marston, 1968: 783) (Young & Duncan, 1994: 8).

Gum dammar (*Agathis dammara* & *Hopis shorea* resin, xylene soluble) = damar (Barbosa, 1974: 86) (Essig) (Hood, 1940: 54) (Lillie, 1953: 59).

Diaphane = juniper gum & phenols, (Batte, 1948: 524) (Hood, 1940: 56) (Lillie, 1953: 64).

Euparal (Colourless not green Euparal vert) = Eucalyptus oil & methyl salicylate, camsal, sandarac, paraldehyde. (Batte, 1948: 524) (Bink, 1979: 160) (Cutler, 1978: 6) (Disney, 1994: 387) (Essig, 1948: 17) (Hangey, 1985: 147) (Herr, 1982: 164) (Hood, 1940: 55) (Knudsen, 1966: 501) (Purvis, 1964: 108) (Ramanna, 1973: 103) (Rattanaarithikul, 1982: 148) (Rawlins, 1992: 56) (Robinson, 1976: 130) (Smithers, 1981: 78) (Steyskal, 1986: 36) (Wagstaffe, 1970: 175) (Wahl, 1989: 181) (Westheide & Purschke, 1988: 152) (Wilkey, 1990: 349) [ASCO Lab and BDH (Merck)].

Lenzol = (Cedar oil), (Gurr, 1963: 88).

Piperine balsam = alkaloid of *Piper nigrum* in chloroform (Frison, 1955: 64).

SYNTHETIC RESINS

Araldite MY753 epoxy resin

(Monniot, 1988: 463) [B & K Resins Ltd].

Aroclor chlorinated diphenyl polymer (Fidiam, 1993: 9) (Frison, 1955: 206) (McLaughlin, 1986: 287) (Wicks, 1946: 121).

B-72 (Morse, 1992: 4).

Caedex (cyclohexanol formaldehyde + plasticiser chlorinated diphenol + xylene) (Lillie, 1953: 69).

Carbowax = (Polyethylene glycol).

Carboxy methyl cellulose CMC = Cellofas & Turtox

(water based)

Methyl cellulose	5 g.
Carbowax (PEG)	2 g.
Diethylene glycol	1 ml.
Ethyl alcohol 95%	25 ml.
Lactic acid	100 ml.
Distilled water	75 ml.

(Barbosa, 1974: 90) (Becker & Heard, 1965: 234) (Beckett, 1982: 97) (Beer, 1954: 1111) (Clark & Morishita, 1950: 789) (Evans, 1961: 82) (Hobson & Banse, 1981: 6) (Holdich & Jones, 1983: 20) (Huryn & Perlmutter, 1988: 432) (Knudsen, 1966: 501) (Krantz, 1978: 89) (Martin, 1977: 109) (New, 1974: 23) (Singer, 1967: 483) (Stehr, 1987: 16) (Walker & Crosby, 1988: 23) [BDH].

Clarite (naphthalene polymer) (Essig, 1948: 19) (Lillie, 1953: 66) (Newell, 1947: 12) (Wicks, 1946: 122).

Clark's **Clearcol** (Chapman, 1985: 117).

Crystalbond 509 thermoplastic cement [Meclec Co.].

Coumarone naphthalene polymer (Frison, 1955: 40) (Lillie, 1953: 67).

DePeX = (Distrene plasticiser xylene) (polystyrene) (Chapman, 1985: 117) (Gagne, 1994: 35) (Lillie, 1953: 65) (Rawlins, 1992: 56) (Westheide & Purschke, 1988: 152) [BDH (Gurr)].

Distrene in xylene + dibutyl phthalate (plasticiser) (polystyrene).

Dymax 304 u.v resin [Intertronics].

Elvacite [BDH].

Elvanol = Polyvinyl alcohol.

Entellan synthetic mounting medium by Merck (Rawlins, 1992: 57).

Epon/Araldite epoxy resin

Dodeceny succinic anhydride	10 ml.
Epon 812	6.2 ml.
Araldite 506	8.1 ml.
Dibutylphthalate	0.75 ml.
2,4,6-tri (dimethyl-aminomethyl phenol)	25 ml.

(Rieger & Ruppert, 1978: 226) (Smith & Tyler, 1984: 261).

Epon epoxy resin

Dodeceny succinic anhydride	100 ml.
Epon 812	62 ml.
+7 parts of solution Nadic methyl anhydrite	89 ml.
Epon 812	100 ml.
+3 parts of solution	
2,4,6-tri (dimethyl-aminomethyl phenol)	1.5%

(Cavey & Cloney, 1973: 150).

Epotek 301 epoxy resin [Intertronics].

Epoxy resins (Loveland: 1986).

Eukitt salicylic acid imbalance caused precipitation of crystals (personal observation K.R.C. Tuck).

Gurr's neutral mounting medium (coumarone and other resins in eucalyptol) = Clearmount (Hockin, 1981: 222) (Huys & Boxshall, 1991: 451) (Lillie, 1953: 68).

Hymount

Hyrax naphthalene resin (Barr, 1973: 16) (Bartsch, 1988: 419) (Beer, 1954: 1111) (Evans, 1961: 82) (Frison, 1955: 64) (McLaughlin, 1986: 287) (Newell, 1947: 7-9) (Singer, 1967: 475) (Wicks, 1946: 121).

Impruv Potting Compound 363 (Loctite) ultra-violet light polymerising plastic (Silverman, 1986: 135).

Impruv Sealant/adhesive 365 (Loctite) ultra-violet light polymerising plastic (Silverman, 1986: 135).

Klearmount epoxy resin (Knudsen, 1966: 501) (Lane, 1974: 7) (Lillie, 1953: 69) (Loveland, 1986: 187).

Lakeside 70C thermoplastic cement (Murray, 1979: 11) [BDH].

Loctite 358 u.v resin [BDH and BSL Ltd].

Lucite (44 & 46) methyl methacrylate plastic (Frison, 1955: 209) (Wicks, 1946: 121).

Malinol (Westheide & Purschke, 1988: 152) (Westheide, 1990: 43).

Meltmount (DeForest, 1987: 154).

Mulford EX-80 polyester resin (Blackburn & Christophel, 1976: 126).

Naphrax

glacial acetic acid	300 ml.
sulphuric acid	100 ml.
Naphthalene	100 g.
Formaldehyde (37% solution)	100 ml.

(Fleming, 1943: 35; 1954: 42) (McLaughlin, 1986: 287) (Ragge, 1955: 8) [NBS].

Novolacs (Union Carbide BRPB 5215) = Phenolic polymer resin (Crumpton, 1980: 347).

Numount (Barton, 1991, 17) (DeForest, 1987: 253) [R.A.Lamb].

Permout (hydrogenated terpene naphthalene polymer) (Herr, 1982: 164) (Knudsen, 1966: 501) (Lillie, 1953: 66) (Stehr, 1987: 16) (Wicks, 1946: 122) [R.A.Lamb].

Petropoxy 154 epoxy resin [BDH and Production Techniques].

Piccolyte (polyterpene polymer resin, xylene soluble) (Frison, 1955: 66) (Herr, 1982: 164) (Knudsen, 1966: 501) (Wicks, 1946: 121).

Pleurax

Phenol crystals	100 g.
sulphur	40 g.
sodium sulphide crystals	2 g.
isopropyl alcohol	100 ml.

(Hepworth, 1994: 21).

Polybed 812 (Ruppert, 1988: 306).

Polyethylene glycol (Carbowax) (PEG) epoxy resin (Walker & Crosby, 1988: 23) (New, 1974: 23).

Polystyrene & Methylene iodide

Polystyrene	15.75 g.
Toluene	50 ml.
Methylene iodide	200 g.

(Czarnecki, 1972: 73).

Polyvinyl acetate

Cellosolve	68%
Diamyl phthalate	12%
Polyvinyl acetate	20%

(Walker & Crosby, 1988) (Ried, 1994)

Polyvinyl alcohol (Danielsson)

PVA Mowiol N 4-982	5 g.
PVA Mowiol N 56-98	5 g.
Ethanol	30 ml.
Lactic acid	105 ml.
Distilled water	105 ml.

(Danielsson, 1985: 383) (Heikinheimo, 1988: 36).

Polyvinyl alcohol (=Elvanol ex DuPont)

Polyvinyl alcohol (low viscosity)	2 g.
70% acetone	7 ml.
Glycerine	5 ml.
Lactic acid	5 ml.

Microscope Slides — Paul A Brown

Distilled water 10 ml.
(Brown, 1951: 263) (Essig, 1948: 18) (Gray, 1950: 290)
(Jones, 1946: 85) (Ribeiro, 1967: 158).

Polyvinyl alcohol Ribeiro

Polyvinyl alcohol 130 g.
Formic acid 40 ml.
Distilled water 50 ml.
(Barbosa, 1974: 96) (Ribeiro, 1967: 159).

Polyvinyl alcohol Ribeiro II

Polyvinyl alcohol 6 g.
Formic acid 10 ml.
Chloral hydrate 72 g.
Phenol 12 g.
(Barbosa, 1974: 97).

Polyvinyl alcohol Salmon

Polyvinyl alcohol (Elvanol 71/24) 5 g.
Lactic acid 10 ml.
glycerine 1 ml.
distilled water 30 ml.
(Salmon, 1954: 66).

Polyvinyl lacto glycerol

Polyvinyl alcohol 16.6g.
Lactic acid 100ml.
Glycerol 10ml.
distilled water 100ml.
(Halliday, 1994: 12)

Polyvinyl lactophenol

Polyvinyl alcohol (Elvonol A) 2.5 g.
Lactic acid 45 ml.
Phenol crystals 45 g.
Distilled water 10 ml.
(Athersuch et al. 1989: 38) (Barbosa, 1974: 98) (Bellinger, 1953: 170) (Bink, 1979: 160) (Henderson, 1990: 26)
(Hodkinson & White, 1979: 3; semi-permanent) (Martin, 1977: 171) (New, 1974: 23) (Nosek, 1973: 84) (Roskam, 1977: 163, [Chroma Gesellschaft Schmid & Co]) (Russell, 1961: 377) (Salmon, 1949: 250) (Wagstaffe, 1970: 176) [Gurr].

Polyvinyl lactophenol Downs

Elvanol PVA solution 56 ml.
Lactic acid 22 ml.
Phenol crystals 22 ml.
(Barbosa, 1974: 96) (Evans, 1961: 81) (Essig, 1948: 19)
(Beer, 1954: 1111) (Downs, 1943: 539) (Lipovsky, 1953: 43).

Polyvinyl lactophenol Jones

Polyvinyl alcohol 6.3 g.
100% alcohol saturated picric acid 18 ml.
Lacto phenol 45 ml.
(Beer, 1954: 1111) (Jones, 1946: 85).

Polyvinyl lactophenol Heinze (used at Harpenden)

Polyvinyl alcohol 10 g.
Chloral hydrate 20 g.
Glycerol 10 ml.
Phenol 1.5% solution 25 ml.
Lactic acid 35 ml.
Distilled water 40-60 ml.
(Heinze, 1953: 177) (MacFarlane, 1991: 80).

Polyvinyl pyrrolidone (PVP) Burstone

Polyvinyl pyrrolidone 50 g.
glycerol 2 g.
Distilled water 50 ml.
(Steedman, 1976: 191).

Pro-Texx (DeForest, 1987: 253).

Realgar (McLaughlin, 1986: 287) (Spence, 1940: 270).
Sira (Wagstaffe, 1970: 176).

Spurr low viscosity embedding medium epoxy resin
Vinylcyclohexane 10 ml.
Diglycidyl ether of polypropyleneglycol 6 ml.
Nonenyl succinic anhydride 26 ml.
(Herr, 1982: 164).

Steedman's post-fixation preservative, PFP

Propylene phenoxetol 5 ml.
Propylene glycol 50 ml.
Distilled water 445 ml.
(Moore, 1979: 494).

Styrax (Fidiam, 1993: 9) (Frison, 1955: 63) (McLaughlin, 1986: 287) (Spence, 1940: 266).

Sucrose benzoate

Sucrose benzoate 60 g.
Polyethylene glycol 600 benzoate 0.25 ml.
Benzyl benzoate 1.75 ml.
Xylene 38 ml.
(Steedman, 1976: 193).

Technicon mounting medium (Westheide & Purschke, 1988: 152).

Tissue-Tek = Polyvinyl alcohol and polyethylene glycol mix (Rawlins, 1992: 51).

Xam maleic polymer terpene resin in xylene + plasticiser (Gurr, 1963: 87) (Lillie, 1953: 69) (DeForest, 1987: 253).

Zeiss W15 (Huys & Boxshall, 1991: 451) (Westheide & Purschke, 1988: 152).

GUM CHLORAL MOUNTANTS (Gray, 1954) (Upton, 1993).

"Andre's" fluid

Gum Arabic 50 g.
Chloral hydrate 125 g.
Glacial acetic acid 50 ml.
Distilled water 50 ml.
(Cunningham, 1972: 906) (Krantz, 1978: 86) (Martin, 1977: 169).

Baylis & Munro

Gum Arabic 15 g.
Chloral hydrate 16 g.
Glucose syrup 10 g.
Glacial acetic acid 5 ml.
Distilled water 20 ml.
(Smart, 1965: 287).

"Berlese" = Imms medium = Lee 1921 = Davidson's

Gum Arabic 15 g.
Chloral hydrate 160 g.
Glucose 10 g.
Acetic acid 5 ml.
Distilled water 20 ml.

(Barbosa, 1974: 93) (Essig, 1948: 14) (Gater, 1929: 367)
(Hood, 1940: 44) (Martin, 1977: 169) (Upton, 1993: 268)

(Wagstaffe, 1970: 175) (Walker & Crosby, 1988: 79)
(Womersley, 1943: 181)[ASCO Lab].

Doetschman

Gum Arabic 20 g.
Chloral hydrate 20 g.
Glycerin 20 ml.
Glucose syrup 3 ml.
Distilled water 35 ml.
(Barbosa, 1974: 94) (Doetschman, 1944: 177) (Upton, 1993: 268).

Eastop & Van Emden

Gum Arabic 48 g.
Chloral hydrate 80 g.
Glacial acetic acid 20 ml.
glucose syrup (50%) 20 ml.
Distilled water 120 ml.
Eastop, 1972: 8) (Upton, 1993: 268).

Ewing

Gum Arabic 20 g.
Chloral hydrate 30 g.
Glycerin 12 ml.
Glucose syrup 3 ml.
Distilled water 35 ml.
(Barbosa, 1974: 96) (Doetschman, 1944: 176) (Upton, 1993: 268).

Faure's

Gum Arabic 30 g.
Chloral hydrate 50 g.
Chloralhydrate of cocaine (optional) 0.5 g.
Glycerine 20 ml.
Distilled water 50 ml.
(Barbosa, 1974: 92) (Bink, 1979: 160) (Evans, 1955: 634)
(Hood, 1940: 46) (Krantz, 1978: 89) (MacFarlane, 1991: 79)
(Martin, 1977: 170) (Morgan & King, 1976: 24) (Upton, 1993: 268) (Wagstaffe, 1970: 176) (Walker & Crosby, 1988: 80).

Foulkes (Foulkes, 1983: 211)

Gum Arabic 16 g.
Chloral hydrate 140 g.
Glycerine 10 ml.
Acetic acid 6 ml.
Distilled water 16 ml.

Higgins' 1983

Gum Arabic (crystals) 30 g.
Chloral hydrate 100 g.
Glycerine 20 g.
Iodine crystals 2 g.
Potassium iodide 1 g.
Distilled water 50 ml.
(Higgins, 1983) (Huys & Boxshall, 1991: 451) (Kristensen & Higgins, 1984: 3) (Westheide & Purschke, 1988: 159).

Hoyer's "125"

Gum Arabic 30 g.
Chloral hydrate 125 g.
Glycerine 20 ml.
Distilled water 50 ml.

(Bartsch, 1988: 419) (Calloway, 1988: 325) (Green & Macquitty, 1987: 15) (Higgins 1983) (Westheide & Purschke, 1988: 152).

Hoyer's

Gum Arabic (crystals) 30 g.
Chloral hydrate 200 g.
Glycerine 20 g.
Distilled water 50 ml.
(Barbosa, 1974: 86) (Beer, 1954: 1111) (Evans, 1955: 635)
(Fain 198X: 169, repairing old mounts) (Gutierrez, 1985: 352) (Jeppson, 1975: 116) (Krantz, 1978: 88) (MacFarlane, 1991: 78) (Martin, 1977: 169) (Pritchard, 1955: 2) (Saito, 1992: 427, temporary) (Soriguer et al., 1993: 113) (Stehr, 1987: 16, questionably permanent) (Upton, 1993: 268) (Wahl, 1984: 228) (Walker & Crosby, 1988: 80).

Jeppson et al.

Gum Arabic 20 g.
Chloral hydrate 125 g.
Glycerine 30 ml.
Sorbitol 30 g.
Distilled water 50 ml.
(Jeppson, 1975: 388) (Upton, 1993: 268).

Puri's

Gum Arabic 8 g.
Chloral hydrate 70 g.
Glycerine 5 ml.
Glacial acetic acid 3 ml.
Distilled water 10 ml.
(Hopkins, 1952: 31) (Kirk & Lewis, 1951: 500) (Smart, 1965: 287) (Upton, 1993: 268).

Reyne

Gum Arabic 60 g.
Chloral hydrate 100 g.
Glycerine 25 g.
Distilled water 100 ml.
(Hamond, 1969: 145) (Reyne, 1950: 40) (Wells, 1988: 383).

Roepke

Gum Arabic 12 g.
Chloral hydrate 20 g.
Glycerine 6 ml.
Distilled water 20 ml.
(Hille Ris Lambers, 1949: 57) (Heie, 1980: 70) (Roepke, 1929: 918).

Singer ("Andre's fluid" ?)

Gum Arabic 50 g.
Chloral hydrate 125 g.
Glycerine 30 ml.
Distilled water 50 ml.
(Bretfield, 1991: 218) (Conde, B.V.N., 1988: 426) (Singer, 1967: 483) (Upton, 1993: 268).

Stroyan (Berlese)

Gum Arabic 12 g.
Chloral hydrate 20 g.
Glucose syrup 5 ml.
Acetic Acid 5 ml.
Distilled water 40 ml.
(Blackman, 1974: 123) (Disney, 1983: 8) (Freeman, 1983: 11) (Henshaw, 1981: 206) (Smith, 1973: 173) (Stroyan, 1949: 6) (Upton, 1993: 268) (Withers, 1989: 12).

Swan's (Berlese)

Gum Arabic 15 g.
Chloral hydrate 60 g.
Glucose syrup 10 g.

Microscope Slides — Paul A Brown

Glacial acetic acid 5 ml.
Distilled water 20 ml.
(Barbosa, 1974: 95) (Nosek, 1973: 84) (Smart, 1965: 287)
(Upton, 1993: 268).

Womersley 1939 (Berlese)

Gum Arabic 12 g.
Chloral hydrate 20 g.
Glucose syrup 4 ml.
Acetic acid 10 ml.
Distilled water 30 ml.
(Womersley, 1939) (Upton, 1993: 268).

Womersley 1943

Gum Arabic 40 g.
Chloral hydrate 50 g.
Phenol 50 g.
Glucose syrup 10 g.
Glacial acetic acid 20 ml.
Distilled water 100 ml.
(Womersley, 1943: 181).

Celochloral

Celodal 1 (Beyer mount) 100 g.
Chloral hydrate 120 g.
Glucose 20 g.
Glacial acetic acid 20 ml.
Distilled water 100 ml.
(Muller, 1961: 70) (Ossiannilsson, 1958: 2).

Keifer's Formaldehyde medium

Gum Arabic 1 g.
Chloral hydrate 14 g.
Glycerine 1 ml.
Sorbitol 3 g.
Formaldehyde 4% solution 5 ml.
Potassium iodide 2 ml.
Iodine 2 ml.
(MacFarlane, 1991: 80).

OTHER AQUEOUS MOUNTANTS

Apathy's fluid

Gum Arabic 50 g.
Sucrose 50 g.
Distilled water 50 ml.
Thymol 0.05 g.
(Grimstone, 1972: 58) (Pantin, 1954: 20) (Steedman, 1976: 189).

Aquamount (Oliver & Meehan, 1993: 17)

[BDH (Gurr) and R.A.Lamb].

Dextrose (Monk, 1938: 174).

Dimethyl hydantoin formaldehyde (DMHF) synthetic resin.

Dimethyl hydantoin formaldehyde 70 g.
Ethyl alcohol 70 ml.
Distilled water 30 ml.
(Bameul, 1990: 233) (Barbosa, 1974: 99) (Cooter, 1991: 57)
(Smith, 1966: 177) (Steedman, 1958: 451; 1976: 189)
[Chemical Intermediates Co Ltd.].

Farrant's medium

Gum Arabic 40 g.
Glycerol 20 g.
Distilled water 40 ml.

(Grimstone, 1972: 58) (Hood, 1940: 45).

Giovacchini's Gelatine

Glycerol 50 ml.
Gelatine 15 g.
Phenol 0.5 g.
Distilled water 55 ml.
(Purvis, 1964: 105).

Glycerine Jelly (Singer)

Glycerine 80 g.
Gelatine 30 g.
Arsenic trioxide saturated solution 150 ml.

(Curry, 1982: 44) (Evans, 1961: 83) (Evens, 1966: 444)
(Garner & Horie, 1984: 93) (Hooper, 1970) (Knudsen, 1966: 501) (Pantin, 1964: 21) (Rawlins, 1992: 56) (Singer, 1967: 483) (Spence, 1940: 143) (Taft, 1983: 399).

Glycerine with Ammonium picrate.

Glycerine Jelly (Kaiser)

Glycerine 100 ml.
Gelatine 15 g.
Phenol 0.25 g.
Distilled water 100 ml.

(Frison, 1955: 169) (Hooper, 1970) (Johansen, 1940: 119)
(Pantin, 1964: 21) (Steedman, 1976: 190).

Glycerine jelly

Glycerine 50 ml.
Gelatine 6 ml.
Phenol crystals 2 g.
Distilled water 42 ml.

(Barbosa, 1974: 92) (Grimstone, 1972: 58).

Glycerine jelly

Glycerine 54g
Gelatine 10g
Phenol 0.5g
Distilled water 50 ml
(Wiles, 1989: 245)

Glycerine, anhydrous

Glycerine 5 ml.
absolute ethyl alcohol 5 ml.
Distilled water 90 ml.

(Chapman, 1985: 113) (Collinson, M.E., 1987: 439)
(Erdtman, 1960: 563) (Green, 1995: 161) (Hood, 1940: 46)
(Hayward, 1985: 30) (Huys & Boxshall, 1991: 451)
(Maybury et al, 1991: 16) (Moore et al., 1991: 45) (Platt & Warwick, 1983: 20) (Pleijel, 1991: 21) (Prescott, 1970: 13)
(Ward, 1944: 200) (Wilson, 1971: 139). [Hopkins & Williams, Gurr (BDH)].

Glycerine/10% formalin (Pleijel, 1991: 21).

Hydramount (Chapman, 1985: 117) (Gurr, 1963: 89).

Karo syrup = dextrose

White Karo corn syrup 10 ml.
Thymol (1% alcohol solution) 1 ml.
Certo 2 ml.
(Knudsen, 1972: 23) (Loveland, 1986: 213) (Taft, 1978: 263).

Keifer's Karo mount

Starch-free white Karo syrup 12 ml.
Chloral hydrate crystals 60 g.
Potassium iodide crystals 0.5 g.

Iodine crystals 2 g.
Formaldehyde solution 10 ml.
(Evans, 1961: 84).

Sodium silicate (water glass) (Bridson & Forman, 1992) (Gerakaris, 1984: 262) (Quisumbing, 1931: 45) (Spence, 1940: 236).

FLUID MOUNTS (Gisin, 1968: 1) (Spence, 1940: 114).

Essig's fluid

Lactic acid 20 parts
Phenol (saturated in distilled water) 2 parts
Glacial acetic acid 4 parts
Distilled water 1 part
(Walker & Crosby, 1988: 80).

Formaldehyde (Diegues & Montero, 1992: 315) (Farrington, 1989: 22).

Formalin Prescott, 1970

Formalin 10 ml.
Ethyl alcohol 95% 30 ml.
Distilled water 60 ml.
(Prescott, 1970: 12).

2% Formalin + 1% cupric acetate solution.

Formalin-acetic acid-alcohol (Prescott, 1970:12) (Wells, 1978).

Gisin's fluid

Lactic acid 179 ml.
Glycerol 36 ml.
Glycerol + saturated picric acid 28 ml.
Formaldehyde (40%) 7 ml.
(Fjellberg, 1980: 9) (Nosek, 1973: 84).

Lactophenol

Phenol 20 ml.
Lactic acid 20 ml.
Glycerine 40 ml.
Distilled water 20 ml.
(Hooper, 1970: 44) (Smart, 1965: 287) (Wells, 1978) [Hopkins & Williams].

Lactophenol

Phenol 30 ml.
Lactic acid 10 ml.
Glycerol 20 ml.
Distilled water 10 ml.
(Huys & Boxshall, 1991: 451).

Monobromonaphthalene (Frison, 1955: 204).

Potassium hydrargiodide (Frison, 1955: 205).

Soft paraffin = Vaseline etc. (Spence, 1940: 115).

Ripart & Petit's fluid

Camphor water 75 ml.
Acetic acid 1 ml.
Cupric acetate 0.3 g.
Cupric chloride 0.3 g.
Distilled water 75 ml.
(Pantin, 1964: 23).

Silicone oil (Chapman, 1985: 118) (Loveland & Centifanto, 1986: 224).

DRY MOUNTS

(Barton, 1991: 17) (Curry, 1982: 42) (Green, 1995: 162) (Hood, 1940: 42).

Black well slides (Athersuch et al. 1989: 39) (Hayward & Ryland, 1979: 34; 1985: 29).

RINGING MOUNTANTS & ADHESIVES

(Disney, 1988: 06) (Gerakaris, 1984: 259) (Gisin, 1968: 1).

APTES = γ -aminopropyl triethoxysilane Sigma tissue section bonding agent. (Rawlins, 1992: 55).

Araldite (Huys & Boxshall, 1991: 451).

Asphalt (Farrington, 1989: 22) (Garner & Horie, 1984: 93) (Johansen, 1940: 120) (Spence, 1940: 141).

Baker's albumin

Aqueous sodium chloride 1% 100 ml.
Sodium p-hydroxybenzoate 0.2 g.
Fresh egg white 100 ml.

Bell's cement = black nitrocellulose lacquer (Garner & Horie, 1984: 93).

Bioseal (Huys & Boxshall, 1991: 451).

Bitumen in toluene (Moore) (Spence, 1940: 141).

Canada balsam (Eastop & van Emden, 1972: 8) (Fjellberg, 1980: 9, unsuitable as it becomes brittle) (Johansen, 1940: 120) (Martin, 1977: 109) (Morgan & King, 1976: 24).

Cellulose nitrate = Celloidin 2-10 % solution in ethyl acetate, plant section adhesive for cryostat, paraffin and glycol methacrylate sections. (Fink, 1987: 98).

Clarite (Evans, 1961: 83).

Cyanoacrylate adhesives (Rawlins, 1992: 51).

Dammar (Garner & Horie, 1984: 98) (Henshaw, 1981: 206).

Eukitt (Fjellberg, 1980: 9).

Euparal (Freeman, 1983: 11) (Gutierrez, 1985: 353) (MacFarlane, 1991: 81) (Martin, 1977: 109).

Glyceel = linseed oil, alcohol, nitrocellulose, butyl acetate & toluol = Zut. (Disney, 1983: 8) (Gutierrez, 1985: 353) (Hooper, 1970: 51) (Huys & Boxshall, 1991: 451) (MacFarlane, 1991: 81) (Martin, 1977: 109).

Glyptal (red alkyd enamel resin insulation paint made by General Electric) (Foulkes, 1983: 211) (MacFarlane, 1991: 81) (Martin, 1977: 109) (Travis, 1968: 24) (Wahl, 1984: 228) (Wu, 1986: 87).

Haupt's adhesive (for plant sections)

Glycerol 15 ml.
Gelatine 1 g.
Phenol crystals 2 g.
Distilled water 100 ml.
(Johansen, 1940: 150) (Purvis, 1964: 105).

Johansen plant section adhesive

Acetone 2 ml.
Methyl benzoate 1 ml.
Distilled water 8 ml.
(Johansen, 1940: 125).

Laktoseal (Morgan & King, 1976: 24).

Mayer's adhesive (for plant sections)

Fresh egg white 50 ml.
Glycerol 50 ml.
Thymol 1 g.

(Johansen, 1940: 125) (Purvis, 1964: 105).

Murrayite

(Eastop & van Emden, 1972: 8) (Essig, 1948: 20) (Hille Ris Lambers, 1949: 57) (Huys & Boxshall, 1991: 451).

Nail varnish thinned with butyl acetate-acetone (typical formulation ex Wells)

Nitrocellulose	10 g.
Resin	10 g.
Plasticiser	5 g.
Ethyl alcohol	10 ml.
Ethyl acetate	20 ml.
Butyl acetate	20 ml.
Toluene	25 ml.

(Disney, 1994: 388) (Fjellberg, 1980: 9) (Wells, 1978).

Norland optical adhesive

Polyisobutylene 5-10% in petroleum ether adhesive for carbowax plant sections. (Fink, 1987: 98).

Polyvinyl Alcohol plant section adhesive

Polyvinyl alcohol	2 g.
Vinyl-triethoxysilane	0.2 g.
Distilled water	100 ml.
Methyl benzoate	0.2 g.

Dilute to 110 to 150 with distilled water for working solution (Fink, 1987: 31).

Shellac (dewaxed in alcohol and plasticised with 1% of castor oil)

(Barton, 1991: 17) (Frison, 1955: 204) (Garner & Horie, 1984: 93) (Gerakaris, 1984: 259) (Gray, 1954: 651) (Johansen, 1940: 120).

Thorn ringing compound thinned with acetone

Nitrocellulose solution	2 parts
Polymerised linseed oil	1 part

(Essig, 1948: 20).

Trycolac (Disney, 1988: 106).

Vaseline (temporary) (Prescott, 1970: 14).

Vinyl-triethoxysilane 0.1% solution, water soluble plant section adhesive. (Fink, 1987: 30).

Paraffin Wax

White vaseline	12 parts
paraffin wax	10 parts
Anhydrous lanolin	10 parts

(Gisin, 1968: 1) (Nosek, 1973: 84).

Xam (Fjellberg, 1980: 9).

STAINS

Stains were not part of the survey but here is a list of a few of the stains known to the author.

Acid Fuchsin (New, 1974: 22) (Saito et al., 1993: 597).

Fast Green (New, 1974: 23).

Lignin Pink (Evans, 1955: 632) (New, 1974: 23).

Chlorazol Black E (New, 1974, 23 This stain requires Polyvinyl lactophenol or Methyl Cellulose / Carbowax unlike the other 3) (Robinson, 1976: 129).

Evans Blue (Saito, 1993: 597).

Aniline Blue (Ramanna, 1973: 103).

Orange G (Lee, 1921) (Gray, 1954)

benzenazo-beta-naphthol-disulphonate of soda, water soluble and acidic.

8. Appendix 2: Refractive Indices

MOUNTING MEDIUM	R.I.	Author
air	1.00	(McLaughlin)
water	1.33	(Rawlins)
Polyvinyl alcohol	1.4	(Heikinheimo)
Glycerine 50% solution	1.4	(Reyne)
Doetschman	1.415	(Gray)
Hoyer	1.419	(Gray)
Methyl cellulose fluid (CMC)	1.428	(Clark)
Glycerol jelly	1.428	(Loveland)
Polyvinyl pyrrolidone (PVP)	1.43	(Steedman)
Diatom silica	1.434	(McLaughlin)
Faure	1.437	(Gray)
Polyvinyl Lactophenol		
HA 1	1.438	(Wagstaffe)
Glycerine jelly	1.443	(Steedman)
Polyvinyl Lactophenol MA 4	1.448	(Wagstaffe)
DMHF (in water)	1.457	(Barneul)
DMHF (in alcohol)	1.466	(Barneul)
Polyvinyl acetate	1.466	(Loveland)
Borosilicate glas (perspex)	1.47	(Heikinheimo)
Karo syrup	1.47	(Bennett)
Swan gum chloral	1.470	(Gray)
Glycerol	1.47	(Rawlins)
Cellulose acetate	1.48-1.50	(Fink)
Euparal	1.48	(Rawlins)
Berlese	1.485	(Gray)
Pro-Texx	1.49	(DeForest)
Formvar	1.49	(Fink)
Histoclear	1.49-1.5	
Entellan	1.49-1.50	(Rawlins)
Cellulose nitrate	1.49-1.51	(Fink)
Polyvinyl alcohol (when dry)	1.491-1.53	(Loveland)
Lucite	1.50	(Bennett)
Silicone	1.50-1.51	(Lillie)
Piccolite	1.50-1.52	(Loveland)
Polyisobutylene	1.50-1.51	(Fink)
Permout	1.51	(Loveland)
Clearmount = neutral mounting medium	1.51	(Gurr)
Immersion oil (Zeiss) crown glass	1.51	(Rawlins)
Zeiss W15	1.51	(Doetschman)
Apathy's	1.515	(Westheide)
Piccolyte	1.52	(Steedman)
Gum dammar	1.52-1.54	(Wicks)
Canada balsam	1.52-1.55	(Lillie)
DePeX	1.52-1.53	(Rawlins)
Xam (terpene resin)	1.529	(Rawlins)
Meltmount	1.52-1.54	(Lillie)
fixed & clear cell constituents	1.539	(DeForest)
DMHF (pure resin)	1.54	(Pantin)
Hydramount	1.54	(Steedman)
Clarite	1.54	(Gurr) ???
Diaphane	1.544	(Loveland)
Sira	1.548	(Loveland)
Styrax	1.55	(Wagstaffe)
Caedax	1.56-1.63	(Loveland)
Sucrose benzoate	1.57	(Loveland)
Coumarone	1.57	(Steedman)
Distrene	1.59	(Frison)
Coumarone + cinnamic aldehyde	1.60	(Wicks)
Novolacs	1.63	(Frison)
Polystyrene	1.65	(Crumpton)
Aroclor	1.60	(Loveland)
Hyrax	1.63	(Wicks)
Aroclor 5442	1.65	(McLaughlin)
Clearax	1.66	(McLaughlin)
Pleurax	1.66	(Gurr)
Piperine balsam	1.67	(Hepworth)
Naphrax (naphthalene formaldehyde)	1.68	(Frison)
Polystyrene methylene iodide	1.71	(Fleming)
Hyrax	1.75	(Czamecki)
Realgar	1.82	(Pantin)
	2.40	(McLaughlin)

N.B. Refractive index changes with temperature of the mountant and with the percentage loss of solvent from it. Canada Balsam as used by biologists dissolved in Xylene is 1.497 and dries to 1.532 (Loveland). Thus a specimen can become less visible after it was initially clear when the slide was first made.

9. Appendix 3: Materials and suppliers

Addresses of suppliers to the Natural History Museum.

Inclusion in this list does not indicate that the NHM endorses these particular suppliers or products.

Aquamount, Canada balsam, Cellofas, DePeX, Elvacite, Eukitt, Flourmount, glycerol jelly, lactophenol, Lakeside 70C, Loctite, Petropoxy, polyvinyl lactophenol, Xam neutral medium.

Entellen, Euparal

BDH Merck

Hunter Boulevard

Magna Park

Lutterworth

Leics. LE17 4XN.

Tel. 0800 223344 Fax 01455 558586

Aquamount, Numount, Permout, Canada balsam

R.A.Lamb

Laboratory Supplies

6 Sunbeam Road

LONDON

NW10 6JL

Araldite 753 & 951

B & K Resins

Unit 2, Ashgrove Estate

Bromley

Kent BR1 4TH

Bisley (15) multidrawer paper storage cabinets

Niceday business supplies

Pondtail Close

Horsham

West Sussex

RH12 5HW

Euparal, Berlese

ASCO Laboratories

52 Levenshulme Road

Gorton

Manchester

M18 7NN

Crystalbond

Meclec Co

5-6 Towerfield Close

Shoeburyness

Essex

SS3 9QP

Dimethyl hidantoin formaldehyde

Chemical Intermediates Co Ltd

Barnfield Industrial Estate

Leek

Staffs ST13 5QG

Dymax 304 & Epotek 301

Intertronics

Unit 9

Station Field Industrial Estate

Banbury Road

Kidlington

Oxon OX5 1JD

Histomount, HistoClear, Hydromount

National Diagnostics (UK) Ltd

Unit 4 Fleet Business Park

Itlings Lane

Hessle, Hull

HU13 9XL

Tel. 01482 646022

Fax. 01482 646013

Hill Unit Cabinets

Stephenson Blake

199 Upper Allen Street

SHEFFIELD

S3 7GW

Tel. 0114 2728325

Fax 0114 2720065

Loctite 358

BSL Ltd

1120 London Road

Norbury

LONDON SW16 4DT

Petropoxy 154

Production Techniques

13 Kings Road

Fleet

Hants GU13 9AV

Plastic slide covers

Preservation Equipment Ltd

Church Road

Shelfanger

Diss

Norfolk

IP22 2DG

Tel 01379 657527

Fax 01379 650582

10. Bibliography

Allen, D.E., 1976. *The naturalist in Britain, a social history*. Allen Lane, London. 292 pp.

Athersuch, J., Horne, D.J. & Whittaker, J.E., 1989. Marine and brackish water Ostracods. *Synopses of the British Fauna (New Series)* **43**: 343 pp.

Bameul, F., 1990. Dimethyl hydantoin formaldehyde un excellent milieu de montage en entomologie. *L'Entomologiste* **46** (5): 233-239.

Barbosa, P., 1974. *Manual of basic techniques in insect histology*. Autumn Publishers, Amherst, Massachusetts. 245 pp.

Barr, D., 1973. Methods for the collection, preservation and study of water mites (Acari Parasitengona) *Royal Ontario Museum, Life Science Miscellaneous Publications* pp 23.

Barton, B., 1991. The collection and mounting of recent Foraminifera. *Balsam Post* **15**: 16-18.

Bartsch, I., 1988. Halacaroidea chapter 44 [in] Higgins, R.P. & Thiel, H., [eds] *Introduction to the study of Meiofauna*. Smithsonian Institution Press, Washington D.C. 488 pp.

Microscope Slides — Paul A Brown

- Batte, E.G.**, 1948. A technique for mounting ectoparasites. *Journal of Economic Entomology* **41** (3): 523-524.
- Baylac, M.**, 1986. Un montage simple des insectes de petite taille dans le baume du Canada. *L'Entomologiste* **42** (5): 311-312.
- Becker, D.A. & Heard, R.G.**, 1965. A permanent fixative — stain — mountant for microscopic monogenetic Trematodes and minute Nematodes. *Turtax News* **43** (9): 234-235.
- Beckett, D.C. & Lewis, P.A.**, 1982. An efficient procedure for slide mounting of larval Chironomids. *Transactions American Microscopical Society* **101** (1): 96-99.
- Beer, R.E.**, A Revision of the Tarsonemidae of the Western Hemisphere. *University of Kansas Science Bulletin* **36** (2): 1091-1387.
- Beirne, B.P.**, 1962. *Collecting, preparing and preserving insects*. Canada Department of Agriculture. 133 pp.
- Bellinger, P.F.**, Salmon's fluid, a new mounting medium for slides of small larvae and larval pelts of Lepidoptera. *The Lepidopterist's News* **7** (5-6): 170-171.
- Bennett, A.H., Jupnik, H., Osterberg, H. & Richards, O.W.**, 1951. Phase Microscopy, principles and applications. Chapman & Hall, London. 320 pp.
- Bink, F.A.**, 1979. Methods for mounting Aleyrodidae specimens. *Entomologische Berichten, Deel* **39**: 158-160.
- Blackburn, D.T. & Christophel, D.C.**, 1976. A method of permanently mounting biological tissue cleared in Herr's four-and-a-half clearing fluid. *Stain Technology* **51** (2): 125-129.
- Blackman, R.L.**, 1974. *Aphids*. Ginn, London. 175 pp.
- Blackman, R.L. & Eastop, V.F.**, 1994. *Aphids on the World's Trees*. CAB International / NHM, Wallingford, Oxon. 987 pp.
- Boorman, J. & Rowland, C.**, 1988. A key to the genera of British Ceratopogonidae (Diptera). *Entomologist's Gazette* **39**: 65-73.
- Bridson, D. & Forman, L.**, 1992. *The herbarium handbook*. Royal Botanic Gardens, Kew, London. 303 pp.
- Brinkhurst, R.O.**, 1982. British and other marine and estuarine Oligochaetes. *Synopses of the British Fauna (New Series)* **21** 127 pp.
- Brown, H.J.**, 1951. Experiences in the use of Polyvinyl alcohol as a mountant. *Microscope* **8**: 263-265.
- Cavey, M.J. & Cloney, R.A.**, 1973. Osmium-fixed and Epon-embedded whole mounts of delicate specimens. *Transactions American Microscopical Society* **92**: 148-151.
- Chapman, J.L.**, 1985. Preservation and durability of stored palynological specimens. *Pollen et Spores* **27** (1): 113-120.
- Clark, E.W. & Morishita, F.**, 1950. C-M medium. A mounting medium for small insects, mites and other whole mounts. *Science NY* **112**: 789-790.
- Clastrier, J.**, 1984. Le montage des petits insectes au baume du Canada. *L'Entomologiste* **40** (4): 175-181.
- Clifford, C.M. & Lewers, D.T.**, 1960. A rapid method for clearing and mounting the genitalia of female Ixodes ticks. *Journal of Parasitology*. **46** (6): 802.
- Coe, R.L.**, 1966. Diptera Pipunculidae. *Handbooks for the Identification of British Insects* **10** (2c): 19-20.
- Collinson, M.E.**, 1987. Special problems in the conservation of palaeobotanical material. *Geological Curator* **4** (7): 439-445.
- Conde, B.V.N.**, 1988. Palpigradida. Chapter 46 [in] Higgins, R.P. & Thiel, H., [eds] *Introduction to the study of Meiofauna*. Smithsonian Institution Press, Washington D.C. 488 pp.
- Cooper, D.W.**, 1988. The preparation of serial sections of platyhelminth parasites, with details of the materials and facilities required. *Systematic Parasitology* **12**: 211-229.
- Cooter, J.**, 1991. Examination of Coleopterous genitalia. [in] *A Coleopterist's handbook*. 3rd edition. Amateur Entomologist's Society, Feltham, Middlesex. pp 53-59.
- Crumpton, W.G. & Wetzel, R.G.**, 1980. Novolacs synthetic resins suitable for mounting biological materials. *Transactions American Microscopical Society* **99** (3): 347-348.
- Cunningham, J.L.**, 1972. A miracle mountant fluid for permanent whole-mounts of microfungi. *Mycologia* **64**: 906-911.
- Curds, C.R.**, 1982. British and other freshwater ciliated Protozoa. *Synopses of the British Fauna (New Series)* **22**: 387 pp.
- Curry, A., Grayson, R.F. & Hosey, G.R.**, 1982. *Under the microscope*. Poole, Dorset, Blandford Press. 160 pp.
- Cutler, D.F.**, 1978. *Applied Plant Anatomy*. London, Longmans 6.
- Czarnecki, D.B., & Harvey, D.W.**, 1972. A new mounting medium for diatoms. *Transactions American Microscopical Society* **91** (1): 73.
- Dales, R.P.**, 1981. *Practical Invertebrate Zoology*. Blackwell, Oxford. 356 pp.
- Danielsson, R.**, 1984. Polyviol as mounting medium for aphids (Homoptera Aphidoidea) and other insects. *Entomologica Scandinavica* **15**: 383-385.
- De Forest, P.R., Shankles, B., Sacher, R.L. & Petraco, N.**, 1987. Meltmount 1.539 as a mounting medium for hair. *Microscope* **35**: 249-265.
- Dieguez, C. & Montero, A.**, 1993. Curatorial and conservation aspects of the E. Caballero Bellido Diatom collection at the Museo Nacional de Ciencias Naturales, Madrid. [in] *International Symposium & First World Congress on preservation and conservation of Natural History collections* **2** 309-321.
- Disney, R.H.L.**, 1983. Scuttle Flies, Diptera Phoridae. *Handbooks for the Identification of British Insects* **10** (6): 7-9.
- Disney, R.H.L.**, 1989. Berlese or Balsam? *Antenna* **13** (2): 48.
- Disney, R.H.L.**, 1994. *Scuttle Flies The Phoridae*. Chapman & Hall, London. 467 pp.

- Disney, R.H.L. & Henshaw, D.H. de C.**, 1988. Berlese fluid for slide mounting insects. *Antenna* **12** (3): 106-107.
- Doetschman, W.H.**, 1944. Some suggestions in microtechnique particularly useful in micro-entomology and parasitology. *Transactions of the American Microscopical Society* **63** (2): 175-178.
- Dolling, W.R.**, 1991. *The Hemiptera*. Natural History Museum, Oxford University Press. London. pp.
- Doncaster, J.P.**, 1961. *Francis Walker's Aphids*. BM(NH), London. pp 165.
- Downs, W.G.**, 1943. Polyvinyl Alcohol a medium for mounting and clearing biological specimens. *Science New York* **97**: 539-540.
- Duckworth, W.D., Genoways, H.H. & Rose, C.L.**, 1993. *Preserving natural science collections Chronicle of our environmental Heritage*. National Institute for the conservation of Cultural Property, Washington. pp 78.
- Eastop, V.F.**, 1984. The acquisition and processing of taxonomic data. [in] *Evolution and Systematics of Aphids. Proceedings of the International Aphidological Symposium at Jablonna*, 1981. Polska Akademia Nauk, Ossilinium, Warsaw, pp. 245-270.
- Eastop, V.F. & van Emden, H.F.**, 1972. The insect material pp 6-9 [in] van Emden, H.F. ed *Aphid Technology*. London, Academic Press. 344 pp.
- Erdtman, G.**, 1960. The Acetolysis method. A revised description. *Svensk Botanisk Tidskrift* **54** (4): 561-564.
- Erdtman, G. & Erdtman, H.**, 1933. The Improvement of pollen-Analysis technique. *Svensk Botanisk Tidskrift* **27** (3): 347-357.
- Essig, E.O.**, 1948. Mounting aphids and other small insects on microscope slides. *Pan Pacific Entomologist* **24** (1): 9-22.
- Evans, G.O. & Browning, E.**, 1955. Techniques for the preparation of mites for study. *Annals & Magazine of Natural History* **8**: 631-635.
- Evans, G.O., Sheals, J.G. & Macfarlane D.**, 1961. *The terrestrial Acari of the British Isles An introduction to their morphology, biology and classification*. Volume 1 Introduction and biology. BM (NH), London. 219 pp.
- Evans, E.D.**, 1961. Some notes on Glycerine mounting media. *Journal Quekett Microscopical Club* **4**: 444-446.
- Fain, A.**, 198X. A method of remounting old preparations of Acarines without raising or displacing the cover slip. *International Journal Acarology* **6** (2): 169-170.
- Farrington, M.**, 1989. Notes on a method of making fluid mounts. *Balsam Post* **11**: 22.
- Fidiam, J.F.**, 1993. Brief history of mounting specimens for the microscope. *Balsam Post* **21**: 5-9.
- Fink, S.**, 1987. Some new methods for affixing sections to glass slides. *Stain Technology* **62**: 27-32, 93-99, 349-354.
- Fjellberg, A.**, 1980. *Identification Keys to Norwegian Collembola*. Norsk Entomologisk Forening. 152 pp.
- Fleming, W.D.**, 1943. Synthetic mounting medium of high refractive index. *Transactions Royal Microscopical Society* **63**: 34-38.
- Fleming, W.D.**, 1954. Naphrax A synthetic mounting medium of high refractive index. New and improved methods of preparation. *Journal Royal Microscopical Society* **74**: 42-44.
- Foulkes, G.D.**, 1983. A new method of mounting Acarines and other meiofauna. *International Journal of Acarology* **9** (4): 211-213.
- Fox, I.**, 1942. The use of creosote in mounting fleas and other arthropods on slides. *Science NY* **96**: 478.
- Freeman, P.**, 1983. Sciarid Flies (Diptera Sciaridae). *Handbook for the Identification of British Insects* **9**: 10-11.
- Freeman, P.**, 1987. Sciarid Flies (Diptera) new species and records with notes on the Tuomikoski collection. *Entomologist's Monthly Magazine* **123**: 195-196.
- Frison, E.**, 1955a. Coumarone Resin as a mounting medium. *Microscope* **10**: 39-42 & 63-68.
- Frison, E.**, 1955b. Some further experiments with synthetic resins as mounting media of high and low refractive indices. *Microscope* **10**: 204-210.
- Gagne, R.J.**, 1994. *The gall midges of the Neotropical region*. Comstock/Cornell University Press, Ithaca & London. 352 pp.
- Garner, R. & Horie, C.V.**, 1984. The conservation and restoration of slides of mosses. *Studies in Conservation*. **29**: 93-99.
- Gatenby, J.B. & Painter, T.S.**, 1937. *The microtome's vade-mecum*. 10th edn. Churchill, London. pp 221.
- Gater, B.A.R.**, 1929. An improved method of mounting Mosquito larvae. *Bulletin of Entomological Research* **19**: 367-368.
- Gates Clarke, J.F.**, 1941. The preparation of slides of the genitalia of Lepidoptera. *Bulletin of the Brooklyn Entomological Society* **36**: 149-161.
- Gerakaris, J.G.**, 1984. Slide ringing revisited. *Microscope* **32**: 259-264.
- Gilbert, P. & Hamilton, C.J.**, 1990. *Entomology a guide to information sources* 2nd edn. Mansell, 259 pp.
- Gill, R.J.**, 1988 The Scale Insects of California Part 1 The Soft Scales (*Homoptera Coccoidea Coccidae*) California Department of Food and Agriculture. *Technological series in Agricultural Systematics and Plant Pathology* **1** 132 pp.
- Gisin, H.**, 1968. A cavity-slide technique for preparing permanent fluid preparations of small organisms. *Revue Ecol. Biol. Sol. T.V.* **4**: 581-583.
- Golden, J.**, 1989. Golden oldies curating SEM specimens. *Collections Forum* **5** (1): 17-26.
- Gooday, A.J.**, 1988. Sarcomastigophora. Chapter 17 [in] Higgins, R.P. & Thiel, H., [eds] *Introduction to the study of Meiofauna*. Smithsonian Institution Press, Washington D.C. 488 pp.
- Goodway, M.**, 1992. Storage system for microscope preparations. [IN] Rose, C.L. & Tones, A.R., eds. *Storage of natural history collections Ideas and practical solutions*. Society for the preservation of natural history collections. York, PA, USA.

- Gray, P.**, 1954. *The Microtome's Formulary and guide*. Constable, London. pp 794.
- Gray, P. & Wess, G.**, 1950. The use of Polyvinyl Alcohol and its derivatives as microscopical mounting media. Part 1 Water miscible mounting media. *Journal of the Royal Microscopical Society* series III **70** (3): 287-291.
- Green, J. & Macquitty, M.**, 1987. Halacarid Mites. *Synopses of the British Fauna (New Series)* **36** 178 pp.
- Green, O.R.**, 1995. Pitfalls, problems and procedures in micropalaeontological preparation and conservation. *Geological Curator* **6**(4): 157-166.
- Grimaldi, D.**, 1993. The care and study of Fossiliferous Amber. *Curator* **36** (1): 31-49.
- Grimstone, A.V. & Skaer, R.J.**, 1972. A guidebook to microscopical methods. Cambridge University Press, London. 134 pp.
- Gurr, E.**, 1956. *A practical manual of medical and biological staining techniques*. Leonard Hill, London. 451 pp.
- Gurr, G.T.**, 1963. *Biological staining methods*. Gurr Ltd, London. 116pp.
- Gutierrez, J.** 1985. Mounting techniques 1.5.3 [in] Helle, W. & Sabelis, eds. *World crop pests; Spider Mites, their biology, natural enemies and control* **1A** Elsevier, Amsterdam. 405 pp.
- Halliday, R.B.**, 1994. Ed Microscope Slide Mounting Media. Results of Informal Survey broadcast over the entomology and systematics electronic bulletin boards.
- Hamond, R.**, 1969. Methods of studying the Copepods. *Journal Quekett Microscopical Club* **31**: 137-149.
- Hangey, G. & Dingley, M.**, 1985. *Biological Museum Methods 2 Plant & Invertebrate techniques*. Academic Press Australia. 323 pp.
- Hardwick, D.F.**, 1950. Preparation of slide mounts of Lepidopterous genitalia. *Canadian Entomologist* **82**: 231-235.
- Harris, R.H.**, 1984. A selective Bibliography on preservation, macro & micro-anatomical techniques in Zoology. *Biological Curator's Group Report No.3*, London. 148 pp.
- Hayward, P.J.**, 1985. Ctenostome Bryozoans. *Synopses of the British Fauna (New Series)* **33**: 169 pp.
- Hayward, P.J. & Ryland, J.S.**, 1979. British Ascophoran Bryozoans. *Synopses of the British Fauna (New Series)* **14**: 312 pp.
- Hayward, P.J. & Ryland, J.S.**, 1985. Cyclostome Bryozoans. *Synopses of the British Fauna (New Series)* **34**: 147 pp.
- Heie, O.E.**, 1980. The Aphidoidea (Hemiptera) of Fennoscandia and Denmark. I Mondaridae, Hormaphididae, Thelaxidae and Pemphigidae. *Fauna Entomologica Scandinavica* **9**: 236 pp.
- Heikinheimo, O.**, 1988. Mounting techniques, Aphid collections, Chapter 8.3 [IN] Minks, A.K. & Harrewijn, P., eds. *World crop pests; Aphids, their biology, natural enemies and control* **2B** Elsevier, Amsterdam. 364 pp.
- Heinze, K.**, 1953. Ein Neues Einbettungsmittel für kleine Insekten, insbesondere für blattläuse. *Transactions 9th International Congress of Entomology*, Amsterdam, 1951. 177-179.
- Heming, B.S.**, 1969. A modified technique for mounting Thysanoptera in Canada Balsam. *Entomological News*. **80**: 323-328.
- Henderson, P.A.**, 1990. Freshwater Ostracods. *Synopses of the British Fauna (New Series)* **40**: 228 pp.
- Henshaw, D.J.de C.**, 1981. Observations on the preparation of Berlese's Fluid. *Entomologist's Monthly Magazine* **116** (3): 206.
- Hepworth, A.**, 1994. Diatom mountant - Pleurax. *Balsam Post* **23**: 21-23.
- Herholdt, E.M.**, 1990. *Natural history collections, their management and value*. Transvaal Museum, Pretoria **1** 172 pp.
- Herr, J.M.**, 1982. An analysis of methods for permanently mounting ovules cleared in four-and-a-half type clearing fluids. *Stain Technology* **57**: (3) 161-169.
- Higgins, R.P. & Thiel, H.**, [eds] 1988. *Introduction to the study of Meiofauna*. Smithsonian Institution Press, Washington. 488 pp.
- Hille Ris Lambers, D.**, 1949. On mounting aphids and other soft-skinned insects. *Entomologische Berichten* **13**: 55-58.
- Hobson, K.D. & Banse, K.**, 1981. Sedentary and archannelid polychaetes of British Columbia and Washington. *Canadian Bulletin of Fisheries and Aquatic Sciences* **209** 1-132.
- Hockin, D.C.**, 1981. Gurr's neutral mounting medium a suitable mountant for the Harpacticoida (Copepoda). *Crustaceana* **40** (2): 222-223.
- Hodkinson, I.D. & White, I.M.**, 1979. Homoptera Psylloidea. *Handbooks for Identification of British Insects*, Royal Entomological Society London 2 (5a) 50pp.
- Holdich, D.M. & Jones, J.A.**, 1983. Tanaids. *Synopses of the British Fauna (New Series)* **27** 98 pp.
- Holloway, J.D., Bradley, J.D. & Carter D.J.**, 1987. *CIE guides to Insects important to man 1 Lepidoptera*. Commonwealth Agricultural Bureau International Institute Entomology, Wallingford, Oxon. 262 pp.
- Hood, J.D.**, 1940. *Microscopical whole mounts of Insects*. Cornell University, 57 pp.
- Hooper, D.H.**, 1970. Handling, fixing staining and mounting nematodes [IN] Southey, J.F. *Laboratory methods for work with plant and soil nematodes*. Ministry of Agriculture, Fisheries and Food, ADAS Reference book **402** HMSO. 202 pp.
- Hopkins, G.H.E.**, 1949. Host-associations of the Lice of mammals. *Proceedings Zoology Society London* **119**: 387-604.
- Hopkins, G.H.E.**, 1952. *Mosquitoes of the Ethiopian Region*, Part 1. British Museum (Natural History) London. 355 pp.

- Humes, A.G. & Gooding, R.U.**, 1964. A method for studying the external anatomy of Copepods. *Crustaceana* **6**: 238-240.
- Huron, A.D. & Perlmutter, D.G.**, 1988. Insecta chapter 47 [in] Higgins, R.P. & Thiel, H., *Introduction to the study of Meiofauna*. Smithsonian Institution Press, Washington D.C. 488 pp.
- Huys, R. & Boxshall, G. A.**, 1991. *Copepod Evolution*. Ray Society No 159, London. 468 pp.
- Iharco, F.A., & Almeida, M.T.**, 1987. Montagem e remontagem de afideos para observacao microscopica. *Boletim da Sociedade Portuguesa de Entomologia* **83**: 111-113. (In Portuguese.)
- Imms, A.D.**, 1929. Some methods of technique applicable to Entomology. *Bulletin of Entomological Research* **20**: 165-171.
- Jeppson, L.R., Keifer, H.H. & Baker, E.W.**, 1975. *Mites Injurious to economic plants*. University of California Press, Berkeley. 116-117.
- Jobling, B.**, 1938. The cardboard cell method for the mounting of insects and their parts. *Proceedings Royal Entomological Society London (A)* **13**: 55-57.
- Johansen, A. J.** 1940. *Plant Microtechnique*. McGraw-Hill New York. pp 523.
- Jones, B.**, 1946. Impregnating Polyvinyl alcohol with picric acid for the simultaneous staining and permanent mounting of Acarina. *Proceedings Royal Entomological Society, London (A)* **21**: 85-86.
- Kearns, C.A. & Inouye, D.W.**, 1993. *Techniques for Pollination Biologists*. University Press Colorado. Pp 583.
- Keyser, D.**, 1988. Ostracoda chapter 35 [in] Higgins, R.P. & Thiel, H., [eds] *Introduction to the study of Meiofauna*. Smithsonian Institution Press, Washington D.C. 488 pp.
- Kirk, R. & Lewis, D.J.**, 1951. The Phlebotomines of the Ethiopian Region. *Transactions Royal Entomological Society London* **102** (8): 383-510.
- Knudsen, J.W.**, 1966. *Biological Techniques collecting, preserving and illustrating plants and animals*. Harper International. 525 pp.
- Knudsen, J.W.**, 1972. *Collecting and preserving plants and animals*. Harper & Row, New York. 320 pp.
- Kozarzhevskaya, E.F.**, 1968. Technique of preparing slides for Coccid (Homoptera Coccoidea) determination. *Entomologicheskoye Obozreniye* **47**: 248-253.
- Kosztarab, M. & Kozar, F.** 1988. *Scale Insects of Central Europe*. Akademiai Kiado, Budapest. 456 pp.
- Krantz, G.W.**, 1986. *A manual of Acarology* 2nd edition. Oregon State University, Corvallis. 509 pp.
- Kristensen, R.M. & Higgins, R.P.**, 1984. Revision of Styraconyx (Tardigrada Halechiniscidae), with descriptions of two new species from Disko Bay, West Greenland. *Smithsonian Contributions to Zoology* **391**: 1-40.
- Lambdin, P.L. & Kosztarab, M.**, 1972. A revision of the seven genera related to Lecanodiaspis (Homoptera Coccoidea Lecanodiaspididae). *Studies on the Morphology and Systematics of Scale Insects* **5**: 7-8.
- Lane J.**, 1974. *The preservation and mounting of insects of medical importance*. World Health Organisation VBC/74.502. 20 pp.
- Lane, R. & Crosskey, R.W.**, 1993. *Medical Insects and Arachnids*. Chapman & Hall, London. 723 pp.
- Lewis, D.J.**, 1973. Phlebotomidae (Sand-flies). 3c pp.173-174 [IN] Smith, K.G.V., ed *Insects and other arthropods of medical importance*. British Museum (Natural History), London. 561 pp.
- Lillie, R.D., Zirkle, C. & Greco, J.P.**, 1953. Final report of the committee on histological mounting media. *Stain Technology* **28**: 57-80.
- Lipovsky L.J.**, 1953. Polyvinyl Alcohol with Lacto-phenol, a mounting and clearing medium for Chigger Mites. *Entomological News* **64**: 42-44.
- Liva, M.**, 1983. New candidates for mounting media. *Microscope* **31**: 231-233.
- Loveland, R.P. & Centifanto, Y.M.**, 1986. Mounting media for microscopy. *Microscope* **34**: 181-241.
- MacFarlane, D.**, 1991. *Acarina*. International Institute of Entomology, London.
- Martin, J.E.H.**, 1977. Collecting, preparing and preserving Insects, Mites and Spiders [IN] *The Insects and Arachnids of Canada* part 1 Canada department of Agriculture, Biosystematics Research Institute, Ottawa. pp 182.
- Martin, J.H.**, 1983. The identification of common Aphid pests of tropical agriculture. *Tropical Pest Management* **29** (4): 395-411.
- Martin, J.H.**, 1987. An identification guide to common Whitefly pest species of the world (Homoptera Aleyrodidae). *Tropical Pest Management* **33** (4): 298-322.
- Maybury, C, Morrison, L. & Stewart, V.**, 1991. The search for a reliable mounting medium for recent 'live' foraminifera. *Journal of Micropalaeontology* **10**(1): 16.
- McLaughlin, R.B.**, 1986. Mounting media [in] Diatom microscopy — some preliminary comments. *Microscope* **34**: 285-291.
- Moniot, F. & C.**, 1988. Chapter 54 Tunicata [in] Higgins, R.P. & Thiel, H., [eds] *Introduction to the study of Meiofauna*. Smithsonian Institution Press, Washington D.C. 488 pp.
- Monk, C.R.**, 1938. An aqueous medium for mounting small objects. *Science, New York* **88**: 174.
- Moore, P.D., Webb, J.A. & Collinson, M.E.**, 1991. *Pollen Analysis*. Blackwell, Oxford. 216 pp.
- Moore, S.J.**, 1979. Restoration of the Quekett Microscope slide collection. *Microscopy* **33** (8): 489-494.
- Morgan, C.I. & King, P.E.**, 1976. British Tardigrades. *Synopses of the British Fauna (New Series)* **9**: 133 pp.
- Morse, J.**, 1992. Insect collection conservation. Reports from ECN meeting Reno. *Insect Collection News* **8**: 1-5.
- Mosely, M.E.**, 1943. The preparation of Insects for the microscope; with special reference to Trichoptera. *The Entomologist* **76**: 227-234 & 241-251.

Microscope Slides — Paul A Brown

- Mound, L.A.**, 1966. A revision of the British Aleyrodidae (Hemiptera Homoptera). *Bulletin British Museum (Natural History), Entomology* **17** (9): 29 pp.
- Mound, L.A.**, 1992. Why collect? Responsibilities and possibilities in a Museum of Natural History. *Insect Collection News* **7**: 8-14.
- Mound, L.A. & Pitkin, B.R.**, 1972. Microscopic whole mounts of Thrips (Thysanoptera). *Entomologist's Gazette* **23**: 121-125.
- Müller, F.P.**, 1961. Celochloral nach Ossiannilsson, ein einschlußmittel für die mikroskopische untersuchung kleiner arthropoden. *Wissenschaftliche Zeitschrift der Universität Rostock* **1**: 69-73.
- Murphy, D.H. & Gisin, H.**, 1959. The preservation and microscopic preparation of Anopheline eggs in a Lacto-Glycerol medium. *Proceedings Royal Entomological Society London (A)* **34**: 171-174.
- Murray, J.W.**, 1979. British nearshore Foraminiferids. *Synopses of the British Fauna (New Series)* **16** 68 pp.
- Museums & Galleries Commission.** 1992. Care of Biological collections. Standards in the Museum 2 55 pp.
- New, T.R.**, 1974. Psocoptera. *Handbooks for Identification of British Insects. Royal Entomological Society, London* **1** (7): 102 pp.
- Newell, I.M.**, 1947. A systematic and ecological study of the Halacaridae of Eastern North America. *Bulletin Bingham Oceanographic collection* **10** (3): 7-9.
- Norris, K.R. & Upton, M.S.**, The collection and preservation of Insects. *Australian Entomological Society, Miscellaneous Publication* **3** pp 33.
- Nosek, J.** 1973. *The European Protura, their taxonomy, ecology and distribution.* Museum d'Histoire Naturelle, Geneva. 345 pp.
- Notton, D. G.**, 1995. B.P. Beirne microscope slides of Ichneumonidae at the Natural History Museum, London. *Biology Curator* issue **2**: 15-17.
- Noyes, J. S.**, 1982. Collecting and preserving chalcid wasps (Hymenoptera Chalcidoidea). *Journal of Natural History* **16**: 315-334.
- Noyes, J. & Polaszek, A.**, 1989. Berlese or Balsam? *Antenna* **13** (1)2.
- Nye, W.P.**, 1947. A simple method of mounting Aphids. *Pan Pacific Entomologist* **23** (2): 73-74.
- Oldroyd, H.**, 1970. *Collecting, Preserving and studying insects.* Hutchinson, London. 336 pp.
- Oliver, P.G. & Meechan, C.J.**, 1993. Woodlice. *Synopses of the British Fauna (New Series)* **49** 136 pp.
- Ossiannilsson, F.**, 1958. "Celochloral" — a new mounting medium for insects. *Entomologisk Tidskrift* **79**: 2-5.
- Palma, R.L.**, 1978. Slide-mounting of Lice a detailed description of the Canada Balsam technique. *New Zealand Entomologist* **6** (4): 432-436.
- Palmer D.**, 1993. Fatal flaw fingers fake fossil fly. *New Scientist*, 13th November, **1994** 4-5.
- Pantin, C.F.A.**, 1964. *Notes on microscopical technique for zoologists.* Cambridge University Press pp 77.
- Perkins, E.J.**, 1956. Preparation of Copepod mounts for taxonomic work and for permanent collections. *Nature, London* **178**: 1075-1076.
- Petersen, A.**, 1959. *Entomological Techniques how to work with insects.* Ohio State University. 435 pp.
- Pitkin, B.R.**, 1988. Lesser Dung Flies (Diptera Sphaeroceridae). *Handbook for the Identification of British Insects* **10** (5e).
- Platt, H.M. & Warwick, R.M.**, 1983. *Freeliving marine Nematodes Part 1 British Enoplids.* Cambridge University Press. 304 pp.
- Pleijel, F. & Dales, R.P.**, 1991. Polychaetes British Phyllodoceoids, Typhloscolecoideans and Tomopteroideans. *Synopses of the British Fauna (New Series)* **45**: 202 pp.
- Postek, M.T.**, 19XX. Mounting cleared Botanical preparations. *Stain Technology* **XX**: 178-179.
- Prescott, G.W.**, 1970. *The freshwater Algae.* Brown, Dubuque, Iowa. 348 pp.
- Pritchard, A.E. & Baker, E.W.**, 1955. A revision of the spider mite family Tetranychidae. *Pacific Coast Entomological Society Memoirs series* **2**: 472 pp.
- Puri, I.M.**, 1931. Larvae of Anopheline Mosquitoes with full description of those of the Indian species. *Indian Medical Research Memoirs* **21**: 1-227.
- Purvis M.J., Collier, D.C. & Walls, D.**, 1964. *Laboratory techniques in Botany.* Butterworths, London. pp 371.
- Quate, L.W. & Steffan, W.A.**, 1966. An alternative method of mounting *Phlebotomus* and other small Diptera. *Journal of Medical Entomology* **3** (2): 126.
- Quisumbing, E.**, 1931. Water Glass as a medium for permanently mounting dissections of Herbarium material. *Torrea* **31**: 45-47.
- Ragge, D.R.**, 1955. *The wing venation of the Orthoptera Saltatoria.* BM(NH), London. pp 159.
- Ramanna, M.S.**, 1973. Euparal as a mounting medium for preserving fluorescence of Aniline Blue in plant material. *Stain Technology* **48** (3): 103-105.
- Rattanarithikul, R.**, 1982. A guide to the genera of Mosquitoes (Diptera Culicidae) of Thailand with illustrated keys, biological notes and preservation and mounting techniques. *Mosquito Systematics* **14** (3): 139-208.
- Rawlins, D.J.**, 1992. *Light Microscopy; an introduction to Biotechniques.* Bios Scientific publishers, Oxford. 143 pp.
- Reid, G.**, 1994. The preparation and preservation of collections [in] Stansfield, G., Mathias, J. & Reid, G. eds. *Manual of Natural History Curatorship*, HMSO. 306pp.
- Remaudiere, G.**, 1992. Une methode simplifiee de montage des aphides et autres petits insectes dans le Baume de Canada. *Revue fr. Ent., (N.S.)* **14** (4): 185-186.
- Reyne, A.**, 1950. Faure's fluid as a medium for microscopic mounts of small insects. *Entomologische Berichten* **13**: 37-42. Dutch with English summary.

- Ribeiro, H.**, 1967. A solidifiable formic acid-PVA solution for transporting, preserving and mounting Mosquito larvae and pupae. *Stain Technology* **42**: 158-159.
- Richards, W.R.**, 1964. A short method for making Balsam mounts of Aphids and Scale insects. *Canadian Entomologist* **96**: 963-966.
- Rieger, R.M. & Ruppert, E.**, 1978. Resin embedments of quantitative Meiofauna samples for ecological and structural studies — description and application. *Marine Biology* **46**: 223-235.
- Rose, C.L.**, 1991. The conservation of natural history collections Addressing preservation concerns and maintaining the integrity of research specimens. [in] Cato, P.S. & Jones, C., eds. *Natural History Museums Directions for Growth*. Texas Tech University Press, Lubbock. 252 pp.
- Robinson, G.S.**, 1976. The preparation of slides of Lepidoptera genitalia with special reference to the Microlepidoptera. *Entomologist's Gazette* **27**: 127-134.
- Roepke, W.**, 1929. A new method of making microscopic Aphid preparations. *Transactions Fourth International Congress of Entomology, Ithaca, 1928* **2**: 917-918.
- Roskam, J.C.**, 1978. Biosystematics of insects living on Birch catkins 1 Gall midges of the genus *Semudobia* Kieffe (Diptera Cecidomyiidae). *Tijdschrift voor Entomologie* **120** (6): 153-197.
- Ruppert, E.E.**, 1988. Gastrotricha Chapter 24 [in] Higgins, R.P. & Thiel, H., [eds] *Introduction to the study of Meiofauna*. Smithsonian Institution Press, Washington D.C. 488 pp.
- Russell, C.R.**, 1961. A simple method of permanently mounting Rotifer Trophi. *Journal Quekett Microscopical Club* **28**: 377-378.
- Russell, S.J.**, 1989. Glass slide preservation of SEM mounted Diatoms. *Diatom Research* **4**(2): 401-402.
- Saito, Y. & Osakabe, M.**, 1992. A new fixation method for preparing Mite specimens for optical and SEM microscopic observation. *Applied Entomology & Zoology* **27** (3): 427-436.
- Saito, Y., Osakabe, M., Sakagami, Y. & Yasui, Y.**, 1993. A method for preparing permanent specimens of Mites with Canada Balsam. *Applied Entomology & Zoology* **28** (4): 593-597.
- Salmon, J.T.**, 1951. Polyvinyl alcohol as a mounting medium in microscopy. *Microscope* **8**: 139-142 + 172-173.
- Salmon, J.T.**, 1954. A new polyvinyl alcohol mounting medium. *Microscope* **10**: 66-67.
- Sandlant, G.R.**, 1978. Slide mounting fresh specimens of small soft-bodied Homoptera (Mealybugs, Scale Insects and Aphids). *New Zealand Entomologist* **6** (4): 430-431.
- Scott, T.L.**, 1951. Wanted — an ideal mounting medium. *Microscope* **8**: 206-210.
- Silverman, M.**, 1986. Light-polymerizing plastics as slide mounting media. *Stain Technology* **61**: (3) 135-137.
- Singer, G.**, 1967. A comparison between different mounting techniques commonly employed in Acarology. *Acarologia* **9** (3): 475-484.
- Smart, J.**, 1965. *A Handbook for the identification of Insects of Medical Importance*. British Museum (Natural History) 303 pp.
- Smit, F.G.A.M.**, 1973. Siphonaptera (Fleas) 7 [in] Smith, K.G.V., [ed] *Insects and other arthropods of medical importance*. British Museum (Natural History), London. 561 pp.
- Smit, F.G.A.M.**, 1957. Siphonaptera. *Handbooks for Identification of British Insects* **1** (16): 94 pp.
- Smith, K.G.V.**, 1989. An introduction to the immature stages of British flies. *Handbook for the Identification of British Insects* **10** (14): 10-11.
- Smith, L.**, 1966. Dimethyl hydantoin formaldehyde resin as a mounting medium for nematodes. *Nematologica* **12**: 177.
- Smith, P.S. & Tyler, S.**, 1984. Serial-sectioning of resin-embedded material for light microscopy recommended techniques for micro-Metazoans. *Mikroskopie (Wein)* **41**: 259-270.
- Smithers, C.**, 1981. *Handbook of Insect Collecting*. David & Charles, London. 120 pp.
- Soriguer, R.C., Fandos, P., Martinez, T., Garcia, E. & Lazo, A.**, 1993. Utilizacion de las colecciones de eodermis vegetales, sus posibles aplicaciones y problemas de conservacion. [in] *International Symposium & First World Congress on preservation and conservation of Natural History collections* **1**: 113-118.
- Spence, D.S.**, 1939-40. Notes on Mounting I - VII. *Microscope* **4**.
- Stansfield, G.**, 1992. Conservation and storage zoological collections. [in] Thompson, J.M.A. ed., *Manual of Curatorship A guide to museum practice*. Butterworth / Heinemann, Oxford. pp 756.
- Steedman, H.F.**, 1958. Dimethyl hydantoin Formaldehyde A new water-soluble resin for use as a mountant medium. *Quarterly Journal of Microscope Science* **99** (4): 451-452.
- Steedman, H.F.**, 1976. *Zooplankton fixation and preservation*. UNESCO Press, Paris. 350 pp.
- Stehr, F.W.**, 1987. Techniques for collecting, rearing, preserving and studying immature insects. [in] Stehr, F.W., *Immature Insects*. Kendal/Hunt Dubuque, Iowa. 754 pp.
- Steyskal, G.C., Murphy, W.L. & Hoover, E.M.**, 1986. *Insects and mites techniques for collection and preservation*. United States Department of Agriculture, Agricultural Research Service, Miscellaneous Publication No. **1443** 103 pp.
- Stroyan, H.L.G.**, 1949. *Aphid techniques for advisory entomologists*. Ministry of Agriculture and Fisheries Conference for advisory entomologists, Occasional Notes No. **8** 1-9.
- Swan, D.C.**, 1936. Berlese's fluid remarks upon its preparation and use as a mounting medium. *Bulletin of Entomological Research* **27**: 389-391.
- Taft, C.E.**, 1978. A mounting medium for fresh water plankton. *Transactions American Microscopical Society* **97** (2): 263-264.

- Taft, C.E.**, 1983. A double coverslip method for preparing permanent Algae slides. *Transactions American Microscopical Society* **102** (4): 399-400.
- Travis, B.V.**, 1968. Glyptal — a useful slide ringing compound. *Journal of Medical Entomology* **5** (1): 24.
- Upton, M.S.**, 1991. *Methods for collecting, preserving and studying insects and allied forms*. Australian Entomological Society Miscellaneous publication No.3 4th edn. Brisbane. 86 pp.
- Upton, M.S.**, 1993. Aqueous gum-chloral slide mounting media an historical review. *Bulletin of Entomological Research* **83**: 267-274.
- Wagstaffe, R. & Fidler, J.H.**, 1970. *The preservation of natural history specimens* 1 Invertebrates. Witherby, London. 205 pp.
- Wahl, D.**, 1984. An improved method for preparing exuviae of parasitic Hymenoptera. *Entomological News* **95** (5): 227-228.
- Wahl, D.**, 1989. Further notes on preparation of exuviae of parasitic Hymenoptera. *Entomological News* **100** (4): 181-182.
- Walker, A.K., & Crosby, T.K.**, 1988. *The preparation and curation of insects*. Department of Scientific and Industrial Research series 163. Auckland, New Zealand. pp 92.
- Ward, T.J.**, 1944. Old Glycerine mounts. *Microscope* **9**: 200.
- Webb, Karen**, 1997. Slide Mounting Diatoms. Plant Collections for non-Botanists Workshop part 2. *Biology Curator* **8**: 10-11.
- Wells, J.**, 1978. The use of nail varnish in microscopy. *Microscopy* **33**: (5)
- Wells, J.B.J.**, 1988. Copepoda chapter 37 [in] Higgins, R.P. & Thiel, H., [eds] *Introduction to the study of Meiofauna*. Smithsonian Institution Press, Washington D.C. 488 pp.
- Welsby, F.W.**, 1955. Experiments with Canada Balsam. *Microscope* **10**: 255-258.
- Westheide, W.**, 1990. Polychaetes Interstitial families. *Synopses of the British Fauna (New Series)* **44**: 152 pp.
- Westheide, W. & Purschke, G.**, 1988. Organism processing chapter 10 146-160 [in] Higgins, R.P. & Thiel, H., [eds] *Introduction to the study of Meiofauna*. Smithsonian Institution Press, Washington. 488 pp.
- White, I.M.**, 1988. Tephritid Flies (Diptera Tephritidae). *Handbook for the Identification of British Insects* **10** (5a).
- White, I.M. & Elson-Harris, M.M.**, 1992. *Fruit Flies of economic significance. Their identification and bionomics*. CAB International, Wallingford, Oxon. 601 pp.
- Wicks, L.F., Carruthers, C. & Ritchey, M.G.**, The Piccolite Resins as microscopic mounting media. *Stain Technology* **21** (4): 121-126.
- Wiles, P.R.**, 1989. Two new species of watermite (Torrenticola: Acari) from South East Asia. *Microscopy* **36**: 245-248.
- Wiles, P.R.**, manuscript. The Hydrachnologists' Handbook. University of Buckingham 17pp.
- Wilkey, R.F.**, 1990. Collection, preservation and microslide mounting 1.5.1 [in] Rosen, D. ed. *World crop pests; Armoured Scale Insects, their biology, natural enemies and control* **4A** Elsevier, Amsterdam. 384 pp.
- Wilson, G.J.**, 1971. A method for the recovery of mounted palynological residues. *Mercian Geologist* **4**: 139-141.
- Wirth, W.W. & Marston, N.**, 1968. A method of mounting small insects on microscope slides in Canada balsam. *Annals of the Entomological Society of America* **61** (3): 783-784.
- Withers, P.**, 1989. Moth Flies (Diptera Psychodidae) *Dipterist's Digest* **4**
- Womersley, H.**, 1943. A modification of Berlese's medium for the microscopic mounting of Acarina and other small arthropods. *Transactions Royal Society of South Australia* **76** (2): 181-182.
- Wu, K.W.**, 1986. Review of the polyethylene bottle application technique for sealing microslide preparations of Mites. *International Journal of Acarology* **12** (2): 87-89.
- Young, D.G. & Duncan, M.A.**, 1994. Guide to the identification and geographic distribution of *Lutzomyia* sandflies in Mexico, the West Indies, Central and South America (Diptera Psychodidae). *Memoirs of the American Entomological Institute* **54** 1-863.
- Young, D.G. & Perkins, P.V.**, 1984. Phlebotomine Sand flies of North America (Diptera Psychodidae). *Mosquito News* **44** (2): 263-304.




Fig. 1 Cobb aluminium double coverslip slide mounts, Aphid collection, Entomology dept, NHM.




Fig. 2 Rothschild perspex well mounts, Heteroptera collection, Entomology dept, NHM.




Fig. 3 Dry mounts, Microlepidoptera wing collection, Entomology dept, NHM




Fig. 4 Dry mount card well slides, Copepod collection, Zoology dept, NHM.




Fig. 5 White and black plastic dry Ostracod mounts, Zoology dept, NHM.




Fig. 6 Electron microscope stubs in card well amounts (& Cobb slide), Ostracod collection, Zoology dept. NHM




Fig. 7 Thick Canada balsam mount with celluloid reinforcement, on non-standard sized slide, Trichoptera collection, Entomology dept. NHM.




Fig. 8 Blackening 'Hoyer's' Diptera mounts made by Lewis, Entomology dept. NHM.




Fig. 9 Diptera slide with multiple cover slips, Entomology dept. NHM.




Fig. 10 Diptera cardboard coverlip mounts, Entomology dept. NHM.




Fig. 11 Slide ringing stage with Euparal and Murrayite ringing media, aphid collection, Entomology dept. NHM.




Fig. 12 Wood and card slide trays in wooden boxes, Entomology dept. NHM.




Fig. 13 Wooden slotted slide boxes (stroyan aphid collection), Entomology dept, NHM.




Fig. 14 Wooden Hill unit with vertical drawers, Thysanoptera collection, Entomology dept. NHM.




Fig. 15 Wooden Hill unit with horizontal drawers, Orthoptera collection, Entomology dept. NHM.




Fig. 16 Metal "Bisley" slide cabinet with vertical drawers, Psyllidae collection, Entomology dept. NHM.




Fig. 17 Liquid formalin mounts, Megoptera collection, Entomology dept. NHM.




Fig. 18 Microscope slide plastic and card envelopes, Aphidoidea collection, Entomology dept. NHM.




Fig. 19 Mixed dry, spirit, slide and host plant gall drawer, Diptera collection, Entomology dept. NHM.




Fig. 20 Canada balsam slides made in 1847, Walker aphid collection and 1898, Pierce Lepidoptera genitalia stained with green ink, Entomology dept. NHM.




Fig. 21 Three Euparal and one Canada balsam mounts, Lepidoptera genitalia collection, Entomology dept. NHM




Fig. 22 insect in amber, slide mounted with Canada balsam, the fossil latrine fly hoax, Paleontology dept. NHM.




Fig. 23 Blackening phenol balsam mounts, Aphidoidea collection, Entomology dept. NHM.




Fig. 24 Euparal descaled wing mount, Butterfly collection, Entomology dept. NHM.




Fig. 25 Naphrax wing mounts, Orthoptera collection, Entomology dept. NHM.




Fig. 26 D.H.R. Lambers slide deterioration, Aphidoidea collection, Entomology dept. NHM.




Fig. 27 Crystallising gum choral Aphidoidea mounts, Entomology dept. NHM.





Fig. 28 Blackening gum choral Aphidoidea mounts, Entomology dept. NHM.



Curriculum Vitae



Paul A. Brown

After being awarded a BSc Honours degree in Forestry & Applied Zoology at the University College of North Wales, Bangor in 1977, employment was found at the Natural History Museum, London in the Lepidoptera section, dealing with pinned specimens and with microscope slide preparations of lepidoptera genitalia using Euparal (1977-1979). Since 1980, he has curated and conserved the Sternorrhyncha (aphid) microscope slide collection, learning techniques in mounting whole specimens with gum chloral and Canada balsam mountants. From 1983 he has been author and assistant author to Dr Roger Blackman of papers on aphid taxonomy and became a Fellow of the Royal Entomological Society of London in 1981 and a Member of the Institute of Biology in 1982. In 1994 Paul was awarded an MA degree in Museum Studies at the Institute of Archaeology, University College, London and was awarded the Diploma of the Museums Association in 1995.

Paul now oversees the slide collections within the Entomology Department and consults and lectures on microscope slides within the NHM and to outside bodies. He organised a day conference on slide mountants at the Natural History Museum sponsored by Merck Ltd in 1996 and lectures on the Imperial College MSc course on Methods in Advanced Taxonomy. He is now playing an active role in Conservation, being on the committee of the Natural Sciences Conservation Group and represents NSCG on the Conservation Forum.

